



Cite this article: Castillo-Morales A, Monzón-Sandoval J, Urrutia AO, Gutiérrez H. 2014 Increased brain size in mammals is associated with size variations in gene families with cell signalling, chemotaxis and immune-related functions. *Proc. R. Soc. B* **281**: 20132428. <http://dx.doi.org/10.1098/rspb.2013.2428>

Received: 17 September 2013

Accepted: 5 November 2013

Subject Areas:

neuroscience, genomics, evolution

Keywords:

encephalization index, brain evolution, gene expression

Authors for correspondence:

Araxi O. Urrutia

e-mail: a.urrutia@bath.ac.uk

Humberto Gutiérrez

e-mail: hgutierrez@lincoln.ac.uk

†These authors contributed equally to this study.

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2013.2428> or via <http://rspb.royalsocietypublishing.org>.

Increased brain size in mammals is associated with size variations in gene families with cell signalling, chemotaxis and immune-related functions

Atahualpa Castillo-Morales^{1,†}, Jimena Monzón-Sandoval^{1,†}, Araxi O. Urrutia¹ and Humberto Gutiérrez²

¹Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, UK

²School of Life Sciences, University of Lincoln, Lincoln LN6 7TS, UK

Genomic determinants underlying increased encephalization across mammalian lineages are unknown. Whole genome comparisons have revealed large and frequent changes in the size of gene families, and it has been proposed that these variations could play a major role in shaping morphological and physiological differences among species. Using a genome-wide comparative approach, we examined changes in gene family size (GFS) and degree of encephalization in 39 fully sequenced mammalian species and found a significant over-representation of GFS variations in line with increased encephalization in mammals. We found that this relationship is not accounted for by known correlates of brain size such as maximum lifespan or body size and is not explained by phylogenetic relatedness. Genes involved in chemotaxis, immune regulation and cell signalling-related functions are significantly over-represented among those gene families most highly correlated with encephalization. Genes within these families are prominently expressed in the human brain, particularly the cortex, and organized in co-expression modules that display distinct temporal patterns of expression in the developing cortex. Our results suggest that changes in GFS associated with encephalization represent an evolutionary response to the specific functional requirements underlying increased brain size in mammals.

1. Introduction

Mammalian species in general tend to have larger brain to body size ratios compared with other vertebrates and in some primate and cetacean species this relationship is particularly pronounced [1]. Large brains represent an evolutionarily costly adaptation as they are metabolically expensive, demand higher parental investment than in species with smaller brains and impose a substantial delay in reproductive age [1–5]. In spite of the cost and adaptive impact of larger brains, the precise nature of genomic changes accounting for variations in encephalization across mammalian species is at present poorly understood [6,7].

Whole-genome sequencing efforts have made it possible to study not just individual variations in specific sequences, but also large-scale differences in gene complements between species. Although overall gene number has changed little over the past 800 million years of metazoan evolution, comparative genomic studies have found large disparities among organisms in the number of copies of genes involved in a variety of cellular and developmental processes, and analyses of gene family evolution have shown that instances of gene family expansion and contraction are frequent [8–12]. In a recent analysis of *Drosophila* species, for instance, large numbers of gains and losses have been described, with over 40% of all gene families differing in size among the analysed species. Importantly, the fact that, in these species, rapid gene family size (GFS) evolution is accentuated

in some functional categories strongly suggests that changes in gene number within gene families may reflect evolutionary responses to specific adaptive demands [10]. In this regard, gene duplication events specifically linked to distinct aspects of vertebrate evolution have been described. Examples include the expansion, during early evolution of the vertebrate lineage, of HOX and PAX gene families which are widely believed to have played a key part in the evolution of many known vertebrate innovations [13,14].

A major goal in evolutionary neurobiology is to understand the molecular changes underlying the extraordinary expansion in brain size observed in mammalian evolution. Whether changes in the number of copies of genes involved in distinct cellular and developmental functions has contributed to shaping the morphological, physiological and metabolic machinery supporting brain evolution in mammalian lineages is not known.

By conducting a genome-wide analysis of 39 fully sequenced mammalian species, we set out to establish whether changes in GFS can be linked to increased encephalization. Our results reveal a proportion of gene families displaying a positive association between GFS and level of encephalization significantly larger than expected by chance. This bias occurs most prominently in families associated with specific biological functions. By examining expression data in human tissues, we further found that gene families displaying the highest association between encephalization and GFS are also statistically enriched in genes that are prominently expressed in the brain, with maximal expression in the cortex and displaying an expression signature distinctly associated with cortical development.

2. Methods

(a) Gene family annotations

Annotated gene families encompassing 39 fully sequenced mammalian genomes were obtained from ENSEMBL [15]. In the context of this annotation, a given gene family constitutes a group of related genes that include both paralogues within the same species and orthologues and paralogues from other species. Any given gene can only be assigned to a single gene family. GFS represents the total number of genes per gene family. In order to maximize the number of families covered in this study (more than 10 000), we included all gene families with members present in no less than six of the 39 mammalian species.

(b) Encephalization index

Because larger species have larger brains, it is necessary to estimate brain mass controlling for the allometric effect of body size. We therefore adopted residuals of a log–log least-squares linear regression of brain mass against body mass as this is the most widely accepted index of encephalization (E_i ; electronic supplementary material, table S1) [16,17]. While direct estimates of the ratio of brain mass to body mass have also been used as an alternative encephalization index [2,18], this measure is known to be poorly related to brain complexity across taxa [16,17]. Accurate estimates of brain residuals based on a sample of 493 mammalian species were kindly provided by Gonzalez-Lagos *et al.* [2].

(c) Correlation coefficients of gene family size and encephalization index

Simple Pearson correlations between E_i and GFS as well as multiple regressions (where maximum lifespan (MLSP) was included

as covariate, see below) were carried out using R-based statistical functions. Numerical randomizations to determine statistical significance were conducted using specially written R-based scripts.

(d) Gene ontology terms analysis

Gene ontology (GO) annotations were obtained from the Gene Ontology database (www.geneontology.org). In this study, a particular GO term was associated with a family whenever that term was linked to any of its members in any species. Only terms found to be linked with more than 50 families were examined.

For each GO category, the average Pearson correlation coefficient was calculated. Statistical significance and expected average Pearson correlation per GO was measured using at least 10 000 equally sized random samples taken from the whole gene family population to directly determine the corresponding p -values. Bonferroni correction was used in all analyses to correct for multiple tests.

Enrichment analysis of GO categories was carried out by counting the number of families assigned to each GO term within the analysed set of gene families. However, any bias in family counts per GO within a set of families could be owing to a bias in the overall density of GO annotation events within that sample. In order to adjust for differences in the density of GO annotations between the test and background samples, we divided the family counts per GO from each sample, by the samples' average number of GO annotations per family. Statistical significance was numerically assessed by obtaining the expected (adjusted) number of families per GO in 10 000 equally sized random samples derived from the overall population of gene families.

(e) Maximum lifespan and partial correlation coefficients

MLSP recorded for each species was obtained from the animal ageing and longevity database (AnAge) [19]. To correct for the potential contribution of MLSP to the association between GFS and E_i , partial correlation coefficients were calculated for each gene family, including MLSP as covariate. The resulting partial coefficient represents the contribution of E_i to the variance in GFS which is not explained by variations in MLSP. Only those gene families displaying a significant partial correlation coefficient ($p < 0.01$) between GFS and E_i were considered further.

(f) Phylogenetic relatedness test

Phylogenetic generalized least-square approach (PGLS) and maximum-likelihood estimation of λ -values were carried out using the CAPER module in R. Because the parameter λ measures the degree to which the phylogeny predicts the pattern of covariance of a given trait across species (where λ -values close to 0 represent no phylogenetic autocorrelation while values close to 1 represent full phylogenetic autocorrelation) [20–22], this approach allows us to obtain a single accurate measure of phylogenetic autocorrelation for each individual gene family. In order to remove the effect of phylogenetic relationships from our analysis, we determined the parameter λ for each of the 713 gene families with significant partial correlation coefficients for E_i and GFS (correcting for MLSP) and eliminated all gene families with a significant phylogenetic interdependence ($p < 0.05$ of $\lambda = 0$, and $p > 0.05$ of $\lambda = 1$). This filtering resulted in 501 gene families on which GO enrichment analyses were subsequently carried out as described above.

(g) Gene expression in human brain

RNA-seq data were obtained for 18 052 genes in a total of 16 human tissues, including brain, derived from the Illumina human body map dataset (ENSEMBL v. 62). Individual genes were

categorized as *prominently expressed* in the brain if their expression level in this tissue was the highest or second highest among all 16 tissues included (top 12.5th percentile). Over-representation was assessed by counting the number of these genes within a given sample. Statistical significance was assessed by comparing this count with those observed in 10 000 equally sized random samples drawn from the wider pool of gene families.

(h) Co-expression network analysis

Weighted gene co-expression network analysis was carried out based on pairwise Pearson correlations between the expression profiles obtained from the BrainSpan database (<http://www.brainspan.org>) for over 21 000 genes. Unsupervised hierarchical clustering was used to detect groups, or modules, of highly co-expressed genes following the method described by Zhang & Horvath [23].

3. Results

(a) Gene family size variations in line with encephalization are over-represented in mammals

In order to assess the relationship between encephalization and GFS variations in mammalian taxa, gene family annotations for 39 fully sequenced mammalian genomes were obtained from ENSEMBL [15]. We included in this study all families with members present in no less than six of the 39 mammalian species (see Methods). This resulted in a total of 12 373 non-overlapping gene families encompassing 595 535 genes, with a mean number of 48.13, and a number of copies per gene family per species ranging from 0 to 448.

E_i for each species was defined as the residual of a log–log least-squares linear regression of brain mass against body mass (see Methods). We obtained correlation coefficients for GFS and E_i for each gene family and the resulting distribution of correlation coefficients showed a distinct shift towards positive values (figure 1a). A Monte Carlo simulation of the expected distribution based on random permutations of GFS values across species revealed that the observed bias is highly significant ($p \ll 0.0001$). In total, we found 8789 families with $r > 0$, representing a shift of 2602 gene families from the negative to the positive tail of the distribution relative to the expected equal number of positively and negatively correlated families ($\chi^2 = 2189.608$, $p \approx 0$; figure 1a, inset). This result demonstrates a highly pronounced over-representation of gene families displaying a positive association between GFS and E_i . This observation is not explained by an overall expansion in gene number across species in line with E_i ($r = 0.251$, $p = 0.127$), but rather by an over-representation of small gene families among those highly associated with encephalization, combined with few larger gene families displaying decreases in size.

We next asked whether the observed enrichment of E_i -related GFS variations was unspecific in terms of the gene populations involved or, alternatively, if this enrichment occurred in gene families specifically associated with certain biological functions. To this end, we used functional GO annotations for ‘biological processes’ and carried out two complementary tests to assess deviations (from random expectations) in the distribution of GO terms associated with gene families displaying a high correlation between GFS and E_i . First, we examined whether there were any significant deviations in the mean correlation coefficient of gene families associated with

individual GO terms (see Methods). Out of all 260 functional categories included, only gene families associated with cell–cell signalling, immune response, chemotaxis, neuropeptide signalling pathways and regulation of immune response displayed a significantly higher than expected average correlation values, between GFS and E_i , after Bonferroni correction (figure 1b). By contrast, no significant bias was observed in functional categories containing families with negative average correlations (not shown).

Second, we measured over-representation of GO terms among the gene families whose GFS variations were most significantly associated with E_i ($r > 0$, $p < 0.05$, $n = 1292$). Among these families, we found that GO terms for immune response, chemotaxis, regulation of immune response, female pregnancy, cell–cell signalling, signal transduction, energy reserve metabolic processes, positive regulation of peptidyl-tyrosine phosphorylation and neuropeptide signalling pathways were significantly over-represented after Bonferroni correction (figure 1c). No GO terms were found to be significantly over-represented among gene families with the highest negative covariance between GFS and E_i (not shown). Taken together, these results show that the observed collective variation in GFS in line with encephalization is not randomly distributed across functional categories but is significantly pronounced in families associated with specific biological functions.

(b) Association between gene family size and encephalization is not explained by lifespan variations

A number of studies on brain evolution have uncovered a robust relationship between relative brain size and lifespan [2,24,25]. In agreement with this, we found a strong association between MLSP and E_i among the species included in this study ($r = 0.7912$, $p < 10^{-8}$). Thus, the observed associations between E_i and GFS could be secondary to an underlying association between MLSP and GFS. Of the 1292 most significantly correlated families ($r > 0$, $p < 0.05$), 927 displayed a stronger association with E_i than with MLSP ($r_{(E_i, \text{GFS})} > r_{(\text{MLSP}, \text{GFS})}$), thereby suggesting a preferential contribution of E_i to the observed bias in the correlation distribution ($\chi^2 = 858.74$, $p = 3.3572e^{-187}$, relative to a random equal distribution of stronger associations). GO enrichment analysis was then repeated including only these 927 families revealing a significant over-representation of gene families associated with immune response, chemotaxis, regulation of immune response, energy reserve metabolic processes, female pregnancy, cell–cell signalling, positive regulation of peptidyl-tyrosine phosphorylation and activation of cysteine-type endopeptidase activity involved in apoptotic processes (figure 1d). It is worth noting that the complementary GO enrichment analysis carried out on gene families with both the most significant association between MLSP and GFS ($r > 0$, $p < 0.05$) and a stronger association with MLSP than E_i ($r_{(\text{MLSP}, \text{GFS})} > r_{(E_i, \text{GFS})}$, $n = 1321$), resulted in no significant enrichment of any GO category. These results shows that enrichment of specific GO terms occurred only among gene families preferentially associated with degree of encephalization, whereas GFS variations potentially associated with increased MLSP showed no significant association with any particular functional category.




Figure 1. Enrichment of gene family size variations in line with increased encephalization in mammals. (a) Histogram showing the distribution of correlation coefficients for GFS and E_i in 12 373 gene families encompassing 39 mammalian genomes. A randomization-based estimation of the expected distribution is represented by the dashed line. Inset: distribution of positive and negative correlations relative to the expected distribution (dashed line). (b) Deviations from random expectations in the mean correlation coefficient of gene families associated with individual GO terms (expressed as $-\log(p)$ -value). Only GO categories with a significant bias are shown. (c) Over-representation of GO terms among gene families most significantly associated with encephalization ($p < 0.05$, $n = 1292$). (d) GO enrichment analysis among the families displaying the most significant correlation with encephalization after removing all families with a stronger association with MLSP than with E_i ($n = 927$). (e) GO-terms enrichment analysis among gene families with the most significant positive partial correlation coefficients for E_i after controlling for the contribution of MLSP in a multiple regression analysis ($n = 713$). (f) GO-terms enrichment analysis among gene families with the most significant positive partial correlation coefficients for E_i with no significant phylogenetic interdependence ($n = 501$). Bonferroni-corrected significance thresholds are indicated with a dashed line. Dark bars indicate common GO terms across all five analyses.

Because MLSP may still partly explain the covariance between GFS and E_i even if the correlation coefficient of GFS with E_i is higher than with MLSP, we used multiple regression analysis to obtain partial correlation coefficients between GFS and E_i after controlling for the contribution of MLSP (see Methods). GO terms enrichment analysis was then carried out only among those gene families with the most significant positive partial correlation coefficients

(partial $r > 0$, $p < 0.05$, $n = 713$). This analysis revealed a significant enrichment of families functionally associated with regulation of immune response, chemotaxis, cell-cell signalling and neuropeptide signalling pathways (figure 1e). These results show that variations in GFS specifically associated with encephalization (i.e. not accounted for by variations in MLSP) are also specifically associated with distinct biological functions.

(c) Phylogenetic relatedness does not explain the observed bias in the distribution of gene families associated with encephalization

For a given gene family, any association between Ei and GFS could be the secondary to existing phylogenetic relationships among the species analysed, as in the absence of any selective forces, closely related species will tend to have both similar degrees of Ei and similar GFS [20,22]. In order to determine the degree to which phylogenetic effects contribute to the observed shift in the correlation distribution, we used a PGLS approach (see Methods) [20,22]. Out of 713 gene families with the most significant positive partial correlation coefficients between Ei and GFS (after correcting for MLSP, see previous analysis), we found a total of 501 gene families for which phylogenetic relationships among species could not account for the covariance between GFS and Ei . Among these families, we observed a significant over-representation of gene families associated with regulation of immune response, cell–cell signalling, energy reserve metabolic processes, female pregnancy and activation of endopeptidase activity involved in apoptosis (figure 1f). These findings demonstrate that the over-representation of specific biological functions among those gene families most strongly associated with higher Ei is neither explained by the known association between MLSP and Ei nor by existing phylogenetic relationships among the species analysed.

(d) Gene families with size increases in line with encephalization show expression signatures consistent with brain functions

To assess whether gene family variations in line with encephalization were directly associated with brain function, we characterized the potential relationship between Ei -associated GFS variations and patterns of gene expression in the human nervous system. For this analysis, we selected the top 501 Ei -associated gene families with both the most significant partial correlation coefficient between Ei and GFS and no significant phylogenetic effects (figure 1f). Using available expression data from the Illumina human body map (see Methods), we looked at the possible over-representation of genes highly expressed in the human brain within the selected 501 gene families. Individual genes were categorized as *prominently expressed* in the brain if their expression level in this tissue was the highest or second highest among all 16 tissues included (top 12.5th percentile). Statistical significance was assessed by comparing with equally sized random samples drawn from the wider pool of gene families (see Methods). This analysis revealed a significant enrichment, within these gene families, of genes prominently expressed in the brain (figure 2a). By contrast, no significant enrichment of genes prominently expressed in the brain was detected among those gene families with the strongest association with MLSP and no significant phylogenetic effects (figure 2a).

Genes involved in cortical development have been shown to display higher variance in expression level during the developmental period of the cerebral cortex compared with adulthood [26]. We therefore looked at the possible representation of genes displaying the highest expression variance during human cortical development relative to adulthood, as defined by Sterner *et al.* [26], within the same 501 gene

families and found a significant enrichment of genes displaying this pattern of expression (figure 2b). By contrast, no significant enrichment of these same genes was observed among the top MLSP-associated gene families (figure 2b).

We next asked whether there was any statistical bias in the relative expression of Ei -associated gene families across different brain regions. Using human brain RNA-seq data from the BrainSpan dataset (see Methods), we obtained the average expression for each gene in the cortex, subcortical regions or cerebellum and split them into three categories according to the region where the highest average expression was found. This analysis revealed a statistically significant enrichment, among those genes contained within the top 501 Ei -correlated gene families, of genes maximally expressed in the cortex (figure 2c). No significant enrichment of genes maximally expressed in subcortical regions was observed among these families. By contrast, genes maximally expressed in the cerebellum were found to be significantly under-represented among the top Ei -correlated gene families. Taken together, these results reveal that gene families displaying the highest association between Ei and GFS are enriched in genes that are prominently expressed in the brain, with maximal expression in the cortex and display an expression signature distinctly associated with cortical development.

In order to characterize further the cortical expression profile of Ei -associated gene families, we used a weighted gene co-expression network analysis approach to identify modules of co-expression among genes contained within the top 501 Ei -correlated gene families. Using human developmental expression data derived from the BrainSpan dataset, we identified 18 modules (figure 3a) associated with distinct temporal patterns of expression. Figure 3b shows the time course of expression of six of these modules summarized by the eigengene associated with each module's co-expression matrix. Some of these modules showed the highest expression levels during the early or late fetal period followed by a progressive decline in expression levels with age. This trend may reverse in some instances in late-adult stages (black module, figure 3b) or show a progressive increase throughout development as illustrated by the yellow module.

4. Discussion

Our results reveal a highly significant over-representation of gene families displaying a positive association between GFS and level of encephalization. This bias occurs most prominently in families associated with specific biological functions. The most robust and consistent bias was observed in gene families associated with cell signalling, immune regulation and chemotaxis.

While chemotaxis and cell signalling functions are known to play central roles in the nervous system, the significance of the observed enrichment of immune system-associated functions among gene families displaying the highest association between GFS and Ei is less clear. In recent years, however, signalling and regulatory mechanisms originally described in the immune system have increasingly been found implicated in key neural-specific roles both in the developing and adult nervous system [27–33]. In addition, in the human cerebral cortex, immune system-related functions have been found to be significantly over-represented among genes displaying higher expression variability in the




Figure 2. Relationship between *Eι*-associated GFS variations and patterns of gene expression in the human nervous system. (a) Over-representation of genes prominently expressed in the human brain (top 12.5th percentile) among the top 501 *Eι*-associated or the top MLSP-associated gene families compared to random expectations. (b) Over-representation of genes displaying the highest expression variance during human cortical development relative to adulthood among the top *Eι*-associated or the top MLSP-associated gene families. (c) Percentage of genes maximally expressed in the cortical (CX), subcortical (SC) and cerebellar (CB) regions respectively. Expected values (mean \pm s.e.m.) were numerically determined using sized-matched random samples of genes drawn from the wider pool of gene families. * $p < 0.01$; ** $p < 0.001$; *** $p < 0.0001$.

developing cerebral cortex than in the adult [26], suggesting a substantial involvement of immune-related signals during cortical development.

Our results, showing a significant over-representation of immune-related functions among *Eι*-associated gene families, support the notion of an underlying and substantial overlap in the regulatory and signalling machinery shared by both the immune and nervous system and in particular during development of the latter.

One possible interpretation is that the observed enrichment of immune-related functions among *Eι*-associated gene families reflects an underlying expansion of immune surveillance in mammals that could be in some way permissive to increased encephalization. While we cannot rule out this possibility, at present, there is little evidence in support of any systematically pronounced and sustained expansion of immune functionalities in mammalian lineages [34]. An alternative interpretation is that signalling and regulatory molecular components that were originally involved in immune-specific functions became gradually recruited by the nervous system in response to the developmental and functional demands of increasingly more complex brains.

The observed association between degree of encephalization and variations of GFS in a large number of gene families is further supported by our finding that *Eι*-associated gene families display a transcriptional signature consistent with

brain-specific functions. Indeed, among the gene families most highly correlated with encephalization with no significant phylogenetic effects, we found a statistically significant enrichment of genes prominently expressed in the brain, strongly indicating that these genes are under comparably higher demand in the nervous system relative to other tissues. When restricting the analysis to the relative expression levels within central nervous system regions, we found that these families are enriched in genes prominently expressed in the cortex, suggesting that *Eι*-correlated changes in GFS may have played a substantial role supporting key aspects of cortical evolution. In this regard, it is worth noting that brain evolution in mammalian lineages is characterized by a disproportional expansion of the brain cortex [35,36]. Analysis of the developmental pattern of expression of these families in the human cortex showed that these genes are organized in co-expression clusters or modules with distinct temporal profiles suggesting a substantial involvement of these families in the developmental organization of the brain.

Genes with the highest degree of connectivity within a module are termed hub genes and are expected to be functionally important within the module. By way of illustration, we examined the turquoise module (figure 3b) and identified a member of a zinc finger gene family (gene family ID: ENSFM00620000999432) as its main hub gene. Interestingly, all but two of the 20 members of this gene family in humans




Figure 3. Temporal patterns of cortical expression of *Ei*-associated gene families. (a) Weighted gene co-expression network analysis was used to detect co-expression modules among genes contained within the top 501 *Ei*-associated gene families using human brain temporal expression data, revealing 18 co-expression modules (coloured). (b) Developmental time course of expression of six representative modules summarized by the level of expression of the eigengene associated with each module's co-expression matrix. Birth point is indicated with a dashed line.

are contained within the same co-expression module. Because genes contained within a co-expression module are thought to be functionally related [23,37], the fact that most members of this zinc finger family are found within the same co-expression module strongly suggests that these genes are functionally related during brain development. We reconstructed the phylogenetic tree of this family and found that the observed pattern is the result of a combination of events of gene loss and gene gain from an original set of four ancestral proteins at the base of the mammalian evolution, overall resulting in a steady increase in the number of gene family members in line with increased level of encephalization ($r = 0.7547$, $p = 2.86 \times 10^{-8}$).

5. Conclusion

In this study, we have found a significant over-representation of GFS variations in line with increased encephalization in

mammals. Importantly, this relationship is not accounted for by known correlates of brain size and is not explained by phylogenetic relatedness. The observed bias occurs most prominently in families preferentially expressed in the brain, in particular the cortex, and significantly associated with distinct biological functions.

Based on our results, we propose that variations in GFS associated with encephalization provided an evolutionary support for the specific cellular, physiological and developmental demands associated with increased brain size in mammals.

Acknowledgements. We thank Leticia Ramirez-Lugo from the National University of Mexico for her technical support and helpful comments from two anonymous reviewers.

Funding statement. This work was supported by CONACyT PhD scholarships to A.C.M. and J.M.S., Dorothy Hodgkin Research Fellowship and Royal Society Research Grant to A.U.O. and University of Lincoln support to H.G.

References

1. Roth G, Dicke U. 2005 Evolution of the brain and intelligence. *Trends Cogn. Sci.* **9**, 250–257. (doi:10.1016/j.tics.2005.03.005)
2. Gonzalez-Lagos C, Sol D, Reader SM. 2010 Large-brained mammals live longer. *J. Evol. Biol.* **23**, 1064–1074. (doi:10.1111/j.1420-9101.2010.01976.x)
3. Isler K, van Schaik C. 2006 Costs of encephalization: the energy trade-off hypothesis tested on birds. *J. Hum. Evol.* **51**, 228–243. (doi:10.1016/j.jhevol.2006.03.006)

4. Leonard WR, Robertson ML, Snodgrass JJ, Kuzawa CW. 2003 Metabolic correlates of hominid brain evolution. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **136**, 5–15. (doi:10.1016/S1095-6433(03)00132-6)
5. Weisbecker V, Goswami A. 2010 Brain size, life history, and metabolism at the marsupial/placental dichotomy. *Proc. Natl Acad. Sci. USA* **107**, 16 216–16 221. (doi:10.1073/pnas.0906486107)
6. Dorus S, Vallender EJ, Evans PD, Anderson JR, Gilbert SL, Mahowald M, Wyckoff GJ, Malcom CM, Lahn BT. 2004 Accelerated evolution of nervous system genes in the origin of *Homo sapiens*. *Cell* **119**, 1027–1040. (doi:10.1016/j.cell.2004.11.040)
7. Shi P, Bakewell MA, Zhang J. 2006 Did brain-specific genes evolve faster in humans than in chimpanzees? *Trends Genet.* **22**, 608–613. (doi:10.1016/j.tig.2006.09.001)
8. Demuth JP, De Bie T, Stajich JE, Cristianini N, Hahn MW. 2006 The evolution of mammalian gene families. *PLoS ONE* **1**, e85. (doi:10.1371/journal.pone.0000085)
9. Fortna A *et al.* 2004 Lineage-specific gene duplication and loss in human and great ape evolution. *PLoS Biol.* **2**, E207. (doi:10.1371/journal.pbio.0020207)
10. Hahn MW, Han MV, Han SG. 2007 Gene family evolution across 12 *Drosophila* genomes. *PLoS Genet.* **3**, e197. (doi:10.1371/journal.pgen.0030197)
11. Hughes AL, Friedman R. 2004 Shedding genomic ballast: extensive parallel loss of ancestral gene families in animals. *J. Mol. Evol.* **59**, 827–833. (doi:10.1007/s00239-004-0115-7)
12. Rubin GM *et al.* 2000 Comparative genomics of the eukaryotes. *Science* **287**, 2204–2215. (doi:10.1126/science.287.5461.2204)
13. Soshnikova N, Dewaele R, Janvier P, Krumlauf R, Duboule D. 2013 Duplications of hox gene clusters and the emergence of vertebrates. *Dev. Biol.* **378**, 194–199. (doi:10.1016/j.ydbio.2013.03.004)
14. Holland LZ, Short S. 2008 Gene duplication, co-option and recruitment during the origin of the vertebrate brain from the invertebrate chordate brain. *Brain Behav. Evol.* **72**, 91–105. (doi:10.1159/000151470)
15. ENSEMBL 2012 <http://www.ensembl.org>. (release 64).
16. Herculano-Houzel S. 2011 Brains matter, bodies maybe not: the case for examining neuron numbers irrespective of body size. *Ann. N.Y. Acad. Sci.* **1225**, 191–199. (doi:10.1111/j.1749-6632.2011.05976.x)
17. Herculano-Houzel S, Collins CE, Wong P, Kaas JH. 2007 Cellular scaling rules for primate brains. *Proc. Natl Acad. Sci. USA* **104**, 3562–3567. (doi:10.1073/pnas.0611396104)
18. Deane RO, Nunn CL, van Schaik CP. 2000 Comparative tests of primate cognition: different scaling methods produce different results. *Brain Behav. Evol.* **55**, 44–52. (doi:10.1159/000006641)
19. Tacutu R, Craig T, Budovsky A, Wuttke D, Lehmann G, Taranukha D, Costa J, Fraifeld VE, de Magalhães JP. 2013 Human ageing genomic resources: integrated databases and tools for the biology and genetics of ageing. *Nucleic Acids Res.* **41**, D1027–D1033. (doi:10.1093/nar/gks1155)
20. Freckleton RP, Harvey PH, Pagel M. 2002 Phylogenetic analysis and comparative data: a test and review of evidence. *Am. Nat.* **160**, 712–726. (doi:10.1086/343873)
21. Garland Jr T, Bennett AF, Rezende EL. 2005 Phylogenetic approaches in comparative physiology. *J. Exp. Biol.* **208**, 3015–3035. (doi:10.1242/jeb.01745)
22. Pagel M. 1999 Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884. (doi:10.1038/44766)
23. Zhang B, Horvath S. 2005 A general framework for weighted gene co-expression network analysis. *Stat. Appl. in Genet. Mol. Biol.* **4**, 1128. (doi:10.2202/1544-6115.1128)
24. Allen JS, Bruss J, Damasio H. 2005 The aging brain: the cognitive reserve hypothesis and hominid evolution. *Am. J. Hum. Biol.* **17**, 673–689. (doi:10.1002/ajhb.20439)
25. Barrickman NL, Bastian ML, Isler K, van Schaik CP. 2008 Life history costs and benefits of encephalization: a comparative test using data from long-term studies of primates in the wild. *J. Hum. Evol.* **54**, 568–590. (doi:10.1016/j.jhevol.2007.08.012)
26. Sterner KN *et al.* 2012 Dynamic gene expression in the human cerebral cortex distinguishes children from adults. *PLoS ONE* **7**, e37714. (doi:10.1371/journal.pone.0037714)
27. Crampton SJ, Collins LM, Toulouse A, Nolan YM, O'Keefe GW. 2012 Exposure of foetal neural progenitor cells to IL-1beta impairs their proliferation and alters their differentiation: a role for maternal inflammation? *J. Neurochem.* **120**, 964–973.
28. Gavalda N, Gutierrez H, Davies AM. 2009 Developmental regulation of sensory neurite growth by the tumor necrosis factor superfamily member light. *J. Neurosci.* **29**, 1599–1607. (doi:10.1523/JNEUROSCI.3566-08.2009)
29. Gutierrez H, Hale VA, Dolcet X, Davies A. 2005 NF-kappaB signalling regulates the growth of neural processes in the developing PNS and CNS. *Development* **132**, 1713–1726. (doi:10.1242/dev.01702)
30. Gutierrez H, O'Keefe GW, Gavalda N, Gallagher D, Davies AM. 2008 Nuclear factor kappaB signaling either stimulates or inhibits neurite growth depending on the phosphorylation status of p65/RelA. *J. Neurosci.* **28**, 8246–8256. (doi:10.1523/JNEUROSCI.1941-08.2008)
31. McKelvey L, Gutierrez H, Nocentini G, Crampton SJ, Davies AM, Riccardi CR, O'Keefe GW. 2012 The intracellular portion of G1TR enhances NGF-promoted neurite growth through an inverse modulation of Erk and NF-kappaB signalling. *Biol. Open* **1**, 1016–1023. (doi:10.1242/bio.20121024)
32. Nolan AM, Nolan YM, O'Keefe GW. 2011 IL-1beta inhibits axonal growth of developing sympathetic neurons. *Mol. Cell Neurosci.* **48**, 142–150. (doi:10.1016/j.mcn.2011.07.003)
33. O'Keefe GW, Gutierrez H, Pandolfi PP, Riccardi C, Davies AM. 2008 NGF-promoted axon growth and target innervation requires G1TRL-G1TR signaling. *Nat. Neurosci.* **11**, 135–142. (doi:10.1038/nn2034)
34. Boehm T. 2012 Evolution of vertebrate immunity. *Curr. Biol.* **22**, R722–R732. (doi:10.1016/j.cub.2012.07.003)
35. Nomura T, Gotoh H, Ono K. 2013 Changes in the regulation of cortical neurogenesis contribute to encephalization during amniote brain evolution. *Nat. Commun.* **4**, 2206.
36. Kaas JH. 2013 The evolution of brains from early mammals to humans. *Wiley Interdiscip. Rev. Cogn. Sci.* **4**, 33–45. (doi:10.1002/wcs.1206)
37. Lee HK, Hsu AK, Sajdak J, Qin J, Pavlidis P. 2004 Coexpression analysis of human genes across many microarray data sets. *Genome Res.* **14**, 1085–1094. (doi:10.1101/gr.1910904)