

Reproduction and nutritional stress are risk factors for Hendra virus infection in little red flying foxes (*Pteropus scapulatus*)

Raina K. Plowright^{1,*}, Hume E. Field², Craig Smith², Anja Divljan³, Carol Palmer⁴, Gary Tabor⁵, Peter Daszak⁶ and Janet E. Foley¹

¹Department of Medicine and Epidemiology, School of Veterinary Medicine, University of California, Davis, CA 95616, USA

²Biosecurity Sciences Laboratory, Department of Primary Industries and Fisheries, Locked Mailbag 4, Moorooka, Queensland 4105, Australia

³Institute of Wildlife Research, University of Sydney, New South Wales 2006, Australia

⁴Biodiversity Conservation Division, Department of Natural Resources, Environment and the Arts, PO Box 496, Palmerston, Northern Territory 0831, Australia

⁵Center for Large Landscape Conservation, Yellowstone to Yukon Conservation Initiative, PO Box 1587, Bozeman, MT 59715, USA

⁶Consortium for Conservation Medicine, 460 West 34th Street, New York, NY 10001, USA

Hendra virus (HeV) is a lethal paramyxovirus which emerged in humans in 1994. Poor understanding of HeV dynamics in *Pteropus* spp. (flying fox or fruit bat) reservoir hosts has limited our ability to determine factors driving its emergence. We initiated a longitudinal field study of HeV in little red flying foxes (LRFF; *Pteropus scapulatus*) and examined individual and population risk factors for infection, to determine probable modes of intraspecific transmission. We also investigated whether seasonal changes in host behaviour, physiology and demography affect host–pathogen dynamics. Data showed that pregnant and lactating females had significantly higher risk of infection, which may explain previously observed temporal associations between HeV outbreaks and flying fox birthing periods. Age-specific seroprevalence curves generated from field data imply that HeV is transmitted horizontally via faeces, urine or saliva. Rapidly declining seroprevalence between two field seasons suggests that immunity wanes faster in LRFF than in other flying fox species, and highlights the potentially critical role of this species in interspecific viral persistence. The highest seroprevalence was observed when animals showed evidence of nutritional stress, suggesting that environmental processes that alter flying fox food sources, such as habitat loss and climate change, may increase HeV infection and transmission. These insights into the ecology of HeV in flying fox populations suggest causal links between anthropogenic environmental change and HeV emergence.

Keywords: Hendra virus; *Henipavirus*; *Pteropus scapulatus*; zoonotic emerging infectious diseases; flying fox

1. INTRODUCTION

The emergence of zoonoses from wildlife is a key threat to global human health (Smolinski *et al.* 2003) and has important implications for conservation (Cohen 2000; Daszak *et al.* 2000; Dobson & Foufopoulos 2001). However, elucidating the ecological conditions that contribute to the emergence of these diseases is a major challenge in disease ecology (Plowright *et al.* 2008). To do this often involves the study of both the behaviour of diseases within hosts and macroecological behaviours of wildlife diseases mediated through large-scale processes, which is a logistical and scientific challenge (Plowright *et al.* 2008).

Since the mid-1990s, five new zoonotic viral diseases (Hendra virus, Nipah virus, Menangle virus, Australian bat lyssavirus and Melaka virus) have emerged from Australasian *Pteropus* species bats (Field *et al.* 2001;

Mackenzie *et al.* 2001, 2003; Chua *et al.* 2007). These viruses can cause significant morbidity and mortality in humans and domestic animals (Eaton *et al.* 2006; Chua *et al.* 2007) and are widely distributed in *Pteropus* hosts (Field 2005; Epstein *et al.* 2006). Nevertheless, little is known about their ecology and epidemiology, including how they are transmitted within *Pteropus* populations or from *Pteropus* hosts to domestic animals. The nomadic lifestyle and complex social structure of *Pteropus* bats, as well as the sporadic nature of bat-borne disease outbreaks, has limited large-scale study on these pathogens and limited progress in understanding their dynamics within bats.

Hendra virus (HeV), a paramyxovirus in the genus *Henipavirus*, was the first of these bat-borne viruses to be recognized. Repeated sporadic spillovers of HeV have occurred in eastern Australia since 1994, resulting in four human cases (two fatal) and more than 30 equine deaths (Murray *et al.* 1995; Field *et al.* 2000, 2007; Daszak *et al.* 2006; Hanna *et al.* 2006). All human cases have involved spillover of infection from flying foxes to

* Author and address for correspondence: Consortium for Conservation Medicine, 460 West 34th Street, New York, NY 10001, USA (plowright@conservationmedicine.org).




Figure 1. Field sites where LRFF were captured in the Northern Territory of Australia: Katherine Gorge National Park; Northern Territory (NT) Rural College; Flora River Nature Park; Elsey National Park; Litchfield National Park; and Katherine River.

pasture-fed horses, which appear to act as amplifier hosts, facilitating transmission to people (Daszak *et al.* 2006). HeV does not cause apparent disease in flying fox reservoir hosts (Williamson *et al.* 1998, 1999). However, the high case fatality rate in humans, and the lack of effective treatment or vaccination, has led to HeV being classified as a biosafety level four (BSL-4) pathogen (Eaton *et al.* 2006).

Flying fox populations are highly spatially structured, often consisting of dense aggregations (camps) of tens of thousands to hundreds of thousands of individuals roosting in canopy vegetation during the day, feeding at night. The little red flying fox (LRFF), *Pteropus scapulatus*, is the only nearly exclusively nectivorous *Pteropus* spp. and their dependence on this patchily distributed, ephemeral food resource (Palmer & Woinarski 1999; Palmer *et al.* 2000) drives a nomadic lifestyle, large population sizes and a wide geographical distribution compared with other Australian flying fox species (Hall & Richards 2000; Birt 2004). Recent modelling work (Plowright 2007) suggests that the large number of LRFF populations and their extensive geographical distribution makes them a key reservoir species. As one of only two species that geographically overlap with all known sites of HeV spillover, little red flying foxes may also play an important role in the emergence of HeV into horses. We conducted our longitudinal study of HeV dynamics in LRFF in the northern range of their distribution (figure 1), where they are least migratory due to unfragmented, relatively abundant and diverse flowering vegetation.

(a) Hypotheses

In this paper, we test the following hypotheses on the transmission and ecology of HeV within flying foxes. Experimental infection suggests that HeV can be transmitted transplacentally (Williamson *et al.* 1999). However, we hypothesize that horizontal transmission is the primary mode of transmission in wild populations, for

two reasons. First, HeV has been detected in urine, rectal swabs and throat swabs of experimentally infected animals (K. Halpin 2003–2004, unpublished data), and, second, the highly social nature of flying fox aggregations makes horizontal transmission via these secretions very likely. We hypothesize that pregnancy plays an important epidemiological role, because pregnancy is associated with increased susceptibility to disease in other mammalian systems (Sheldon & Verhulst 1996) and most HeV spillover events coincide with peak flying fox birthing seasons (Field *et al.* 2001, 2007). Lastly, we hypothesize that seasonal increase in contact rates, and the immunosuppressive effects of testosterone associated with mating, may drive temporal fluctuations in HeV dynamics and the risk of spillover to horses and humans. To test these hypotheses, we constructed age-specific seroprevalence curves to examine temporal patterns of exposure. We also sampled flying foxes during each stage of the highly seasonal reproductive cycle (mating, birthing, lactation and weaning) and examined the relative risk of exposure in relation to sex and reproductive status in individuals, as well as seasonal population-level seroprevalence patterns.

2. MATERIAL AND METHODS

(a) Source of the data

A total of 664 LRFF were sampled over five field seasons (November 2004 to March 2006) in the Northern Territory of Australia (figure 1; table 1). Flying fox campsite locations shifted between seasons as the distribution of ephemeral nectar resources changed, requiring sampling at multiple locations within the study area. Sampling periods were chosen to coincide with mating (spring), late pregnancy/early birthing (autumn) and late lactation/early weaning (winter). Sampling was not possible in summer, due to the inaccessibility of sites during the wet season, or in winter 2006, when no LRFF could be located in the study area.

Table 1. Description of risk factors for Hendra virus serostatus included in univariate analysis. (Data were collected from 664 LRFF sampled over five field seasons between November 2004 and March 2006.)

variable	description	no. of animals
sex		
	male	321
	female	343
age		
0–1 month	a pup that is dependent on its mother, usually up to the age of three months; as we were sampling during the peak birthing time, these animals were all less than a month old	8
3 months	juveniles sampled in July; the pup is too large to be carried by the mother, but has not yet reached full size; mandibular third molars have not fully erupted; permanent dentition still fragile with slight pinkish tone (appears almost transparent in places)	50
6 months	juveniles sampled in November; the pup is independent, but has not yet reached full size; all teeth have erupted and have reached their full length; teeth show no signs of wear (cusps are sharp), but slight (diet dependent) lateral staining is possible	64
pre-breeding	an animal that is full size but not sexually mature; from 1 to 2 years; in females, nipples are small and surrounded by fur; males have underdeveloped testes regardless of the season; dentition likely to show signs of staining on lateral surfaces (diet dependent), cusps of premolar teeth are becoming slightly oblong, but there is no wear or staining on canine teeth	198
mature	a full-sized sexually mature animal (potentially breeding); from 2 years to at least 16 years old; mature males have well developed testes (particularly in the mating season); in females that have bred, nipples are enlarged, irrespective of whether they have lactated recently; also, lactating and post-lactating females have less fur around the nipple; teeth are mostly heavily stained from all sides, often showing signs of discoloration and decay, but in various stages of wear	344
season		
spring 2004	November 2004, Elsey National Park	170
autumn 2005	April 2005, Flora Nature Park	103
winter 2005	July 2005, Flora Nature Park ($n=69$) and Northern Territory Rural College ($n=116$)	185
spring 2005	November 2005, Elsey National Park ($n=93$) and Katherine River ($n=87$)	180
autumn 2006	March 2006, Katherine Gorge National Park ($n=18$) and Litchfield National Park ($n=8$)	26
weight	body weight (g)	664
forearm	length from the styloid process of the third metacarpal to the olecranon process of the ulna (mm) when the wing is folded; forearm length does not increase once the animal reaches maturity (approx. 2 years)	664
weight/forearm ratio	body weight (g) divided by forearm length (mm) gives an indication of body condition by taking into account body size	664
lactating	enlarged nipples expressing milk	22
pup	dependent pup attached to its mother	8
pregnant	pregnancy detected via palpation	34

(b) Capture and sampling methods

Flying foxes were caught in mist-nets at 'fly in', the two hours before dawn when animals return to roosting sites after foraging. Mist-nets were raised between either ground-mounted masts, masts placed at either end of a small outboard-powered boat, or hoisted between ropes in the tree canopy. Each animal was anaesthetized via mask induction of isoflurane as described by Jonsson *et al.* (2004), or with an intramuscular injection of ketamine ($6\text{--}7\text{ mg kg}^{-1}$) and medetomidine ($60\text{--}70\text{ }\mu\text{g kg}^{-1}$) followed by reversal with atipamezole ($300\text{--}350\text{ }\mu\text{g kg}^{-1}$).

Sex, reproductive status, forearm length, weight and age were recorded for individual animals. Criteria for classification into age classes are described in detail in table 1. An approximate age for mature flying foxes was determined by tooth wear. We aged a subset of individuals ($n=163$) by the analysis of annual cementum growth layers in an extracted first left mandibular premolar as described by Divljan *et al.* (2006).

Blood was collected by venepuncture from the propatagial (leading edge wing membrane) vein, and in accordance with the Australian Bird and Bat Banding Scheme, each individual was fitted with an individually numbered thumb

band so that recaptured individuals could be identified. After full recovery from anaesthesia, flying foxes were released at the capture site.

Population estimates were not possible due to thickly vegetated terrain, lack of topographic variation and 'fly outs' of large numbers of animals that occurred over a 360° radius. Roughly, we estimated camp sizes to be in the hundreds of thousands of individuals, except for autumn 2006, when only a few thousand individuals could be located.

(c) Serologic tests

Serum neutralization tests (SNTs), the reference standard for HeV (Daniels *et al.* 2001), were performed on serum or plasma samples from each bat in the Biosecurity level 4 laboratory at the Australian Animal Health Laboratory in Geelong, Australia. Neutralization of HeV by the sample at a dilution of 1 : 5 or greater was regarded as a positive test based on titres generated by experimentally infected flying foxes (Williamson *et al.* 1998; Halpin *et al.*, unpublished data). Samples that were unable to neutralize HeV at any dilution were regarded as negative, while samples that were toxic to the cell layer were regarded as unknown. The samples were retested using a Luminex assay (Bossart *et al.*

Table 2. Univariate analysis of risk factors for Hendra virus serostatus in LRFF: proportion seropositive, odds ratios with 95% CIs and chi-square *p*-values for variables included in the analysis.

variable	category	proportion seropositive	odds ratio	95% CI	chi-square <i>p</i> -value
sex and reproductive status	non-pregnant female	0.16	1.00		
	pregnant female	0.83	25.47	9.23–70.22	<0.001
	lactating female	0.52	5.6	2.24–14.01	<0.001
	male	0.23	1.55	1.02–2.38	0.041
age	0–1 month	1.00	3×10^8	0–inf	0.971
	3 months	0.14	9.47	1.1–81.79	0.041
	6 months	0.02	1.00		
	pre-breeding	0.12	8.46	1.12–64.17	0.039
	adult	0.35	32.62	4.46–238.54	<0.001
season	spring 2004	0.31	1.00		
	autumn 2005	0.47	1.66	0.97–2.83	0.066
	winter 2005	0.16	0.41	0.24–0.71	0.002
	spring 2005	0.06	0.14	0.07–0.28	<0.001
	autumn 2006	0.76	6.99	2.63–18.62	<0.001
forearm weight	forearm length (mm)		1.09	1.06–1.12	<0.001
	body weight (g)		1.01	1.00–1.01	<0.001
weight/forearm ratio ^a	body weight (g)/forearm length (mm)		1.75	0.80–3.83	0.163

^a Excluding pregnant females.

2007) to ensure that laboratory error was not a cause of the patterns observed. As HeV does not cause disease-induced mortality in flying foxes (Williamson *et al.* 1998, 1999), and induces long-lived detectable immunity in *Pteropus conspicillatus* and *Pteropus poliocephalus* (spectacled and black flying foxes; Field 2005; H. Field 1998–2001, unpublished data), we assumed that seroprevalence measures cumulative past exposure to HeV.

(d) Statistical approach

All variables (described in table 1) were first screened using a univariate analysis and a chi-square test to check for statistically significant associations with serological status. Data were then analysed using logistic regression and chi-square statistics in R (R Development Core Team 2006). Variables were selected for inclusion in the multivariate model based on the likelihood ratio test using forward and backward methods. Continuous variables were categorized and checked graphically to verify the linearity of the log odds. Potential confounding variables were identified based on a change of greater than 10% in the odds ratio of other variables after adding the potential confounder to the model, previous knowledge and biological reasoning. Adjusted odds ratios and 95% CIs were used to assess the strength of association between risk factors and HeV serostatus.

3. RESULTS

Of the 664 flying foxes sampled over the five field seasons, we obtained 601 SNT results with an overall seroprevalence of 23.62%. Based on univariate analyses, season, age, forearm length, weight, pregnancy, lactation and sex were significantly ($p < 0.05$) associated with serological status (table 2). No individual flying foxes were recaptured.

(a) Stratified analysis

(i) Age and size

Apparent maternal immunity in pups waned over the first six months of life, after which seroprevalence increased

with age, creating a typical concave-shaped curve (figure 2a). Seroprevalence increased with forearm length (figure 2b), a trend that was significant after we accounted for age ($p < 0.001$). The analysis of extracted teeth demonstrated that seroprevalence continues to increase with age within the mature class; however, most teeth were removed during winter 2005 and spring 2005 when there were very few seropositive animals, hence the strength of association between age and serostatus is dampened in this subset of data. Figure 2c illustrates the age distribution of animals from which teeth were removed for analysis.

(ii) Sex and reproductive status

Seroprevalence in males and females differed significantly only during the 2005 birthing season (autumn), when the seroprevalence in pregnant and lactating females was more than four times the seroprevalence in adult males ($p < 0.0001$) and non-reproductive females ($p < 0.0001$; figure 3a). In autumn 2006, no females were pregnant or lactating, presumably due to poor body conditions and nutritional stress observed, yet both males and females of all age groups had a high seroprevalence, equivalent to that of the reproductive females in the 2005 birthing season.

(iii) Season and location within seasons

Seroprevalence differed significantly between seasons (table 2; figure 3b), although the patterns were not annually predictable. Overall seroprevalence in spring 2004 (mating), autumn 2005 (birthing) and autumn 2006 (birthing season but animals not reproducing) was more than twice as high as seroprevalence in winter 2005 (weaning) and spring 2005 (mating). Sampling location within a season was not a significant risk factor for serostatus (table 2): there were no significant differences between the Flora River and the Northern Territory Rural College in winter 2005 ($p = 0.91$), the Katherine River and the Elsey National Park in spring 2005 ($p = 0.10$), or the Litchfield National Park and the Katherine Gorge in




Figure 2. Patterns of HeV seroprevalence in relation to age and size. (a) Age-specific HeV seroprevalence as determined by criteria in table 1: animals less than one month old were all seropositive ($n=8$), reflecting the serostatus of their mothers, which were also all seropositive. Error bars represent 95% binomial CIs. (b) Relationship between forearm length (for animals with forearm length ≥ 124 mm) and HeV seroprevalence. The risk of HeV infection increases with body size. Animals with a forearm length of above 130 mm were all mature animals. Error bars represent 95% binomial CIs. A, 125–130; B, 131–135; C, 136–140; D, 141–145; E, 146–150; F, 151–155; G, 156–160. (c) Age distribution of animals from which teeth were analysed.

autumn 2006 ($p=0.85$). Repeated analyses using luminex test results revealed the same patterns in the data.

(b) Logistic regression analysis

Age, season and sex/reproductive status were included in the main effects model (table 3). In the final model, mature flying foxes, pregnant and lactating females and flying foxes caught in autumn 2006 had a significantly higher risk of being seropositive ($p<0.05$), while flying foxes caught in winter and spring 2005 had a significantly lower risk of being seropositive ($p<0.01$). Notably, when adjusted for sex and age, the odds of being seropositive in autumn 2006 were 14–42 times higher than the odds of being seropositive in any other season. Lactating females were sampled during the birthing season, when seroprevalence was high, and during winter, when seroprevalence was low. In contrast, pregnant females were only sampled




Figure 3. HeV seroprevalence as a function of reproductive status and season. (a) Seroprevalence across reproductive classes in autumn 2005 (preg, pregnant; lact, lactating; not preg/lact, not pregnant or lactating). (b) Overall seroprevalence across seasons (Spr04, spring 2004; Aut05, autumn 2005; Win05, winter 2005; Spr05, spring 2005; Aut06, autumn 2006). Error bars represent 95% binomial CIs.

during birthing; therefore, lactation was associated with a lower relative risk than pregnancy. Forearm length, weight and weight/forearm ratio did not significantly improve the fit of the model after adjusting for other covariates (based on the likelihood ratio test), and were therefore excluded. Two-way interactions among the main effects could not be assessed due to the lack of data in some cells.

4. DISCUSSION

Our analysis provides empirical evidence that nutritional status and reproduction are important drivers of HeV seroprevalence. These results have implications for the management and control of HeV outbreaks, because they indicate that seasonality and environmental stresses may lead to epidemics of HeV in flying foxes. Age-specific seroprevalence curves indicate that horizontal transmission is the principal form of intraspecific transmission, suggesting that contaminated faeces, urine or saliva may be responsible for spillover to other species. Waning acquired immunity in LRFF is suggested, and it could be an important driver of within-host dynamics in this system, as well as in the persistence of HeV within interspecific flying fox communities.

Table 3. Logistic regression model of risk factors for Hendra virus serostatus in LRFF: coefficients, chi-square *p*-values and odds ratios with 95% CIs for variables included in the model.

variable	category	coefficient estimate	chi-square <i>p</i> -value	odds ratio	95% CI
sex and reproductive status	non-pregnant female	reference			
	pregnant female	2.74	<0.001	15.48	4.41–54.35
	lactating female	1.38	0.010	3.96	1.4–11.24
	male	0.02	0.932	1.02	0.61–1.71
age	age 0	19.33	0.972	2 × 10 ⁸	0–inf
	3 months (weaning)	2.19	0.060	8.91	0.91–87.17
	6 months	reference			
	pre-breeding	1.65	0.120	5.19	0.65–41.38
season	mature	2.65	0.011	14.14	1.86–107.75
	spring 2004	reference			
	autumn 2005	–0.70	0.075	0.49	0.23–1.07
	winter 2005	–0.97	0.007	0.38	0.19–0.76
	spring 2005	–1.85	<0.001	0.16	0.07–0.33
	autumn 2006	1.90	<0.001	6.67	2.43–18.31

When experimentally infected with HeV, flying foxes develop viraemia and excrete virus in urine, faeces and saliva for approximately one week (Halpin *et al.*, unpublished data). In their typically dense, three-dimensional aggregations, flying foxes excrete urine and faeces throughout the day, and it is not uncommon to see a fine mist of urine in the camp. Under these conditions, particularly given regular grooming behaviour, horizontal transmission is very likely. The concave curve in figure 2*a* suggests waning maternal immunity followed by cumulative risk of exposure with age and is consistent with horizontal transmission. The finding that maternal antibody status declines over the first six months of life (descending portion of curve in figure 2*a*) corresponds with the findings from Field (2005), in which maternal antibodies in captive-born *P. conspicillatus* and *P. poliocephalus* declined over seven months.

Vertical transmission has been documented in flying foxes on two occasions: in parenterally inoculated pregnant *P. poliocephalus* (Williamson *et al.* 1999; but not in oronasally inoculated pregnant *Pteropus alecto*, Halpin *et al.*, unpublished data) and in two injured free-living flying foxes (*P. alecto* and *P. poliocephalus*) in temporary captivity (Halpin *et al.* 2000). The lack of antibody status in juvenile flying foxes after maternal immunity has waned, along with the concave age-specific seroprevalence curve, suggests that if vertical transmission occurs in LRFF, it is not the predominant form of transmission.

HeV seroprevalence showed a scaling relationship with increasing forearm length, even after we controlled for age (figure 2*b*). This phenomenon suggests the possibility that risk of disease exposure increases with host body size, since forearm length, which does not increase in mature animals, serves as a proxy for adult body size. One explanation for this pattern is that size is associated with dominance rank and that dominant animals have higher contact rates. Behavioural studies of flying foxes have documented hierarchies where dominant males control larger harems of females during mating (Markus 2002), and dominant individuals defend feeding sites against the intrusion of other individuals (Birt 2004).

The fourfold higher seroprevalence in pregnant and lactating females in autumn 2005, compared with seroprevalence in males and pre-breeding females,

indicates that pregnancy and lactation are risk factors for viral infection. Increased susceptibility to infectious disease during pregnancy and lactation has been demonstrated in other mammalian systems (Lloyd 1983; Alexander & Stimson 1988; Festa-Bianchet 1989; Sheldon & Verhulst 1996; Nelson 2004; Cattadori *et al.* 2005; Trevejo *et al.* 2005) due to hormonal suppression of immunity (Alexander & Stimson 1988) and the allocation of resources to reproduction at the expense of immune defence (Sheldon & Verhulst 1996; Nelson 2004). The modified behaviour of pregnant and lactating flying foxes may also promote exposure.

This is the first study to show an association between birthing and HeV infection in a flying fox population, and supports the hypothesis that HeV spillover is linked to flying fox reproduction. A temporal association between the peak birthing season in flying foxes and some disease events in horses (Young *et al.* 1997; Field *et al.* 2000, 2007; Halpin *et al.* 2000), and evidence of vertical transmission (Williamson *et al.* 1999; Halpin *et al.* 2000), has led to speculation that pasture contaminated with aborted fetuses or birthing fluids was the source of infection for horses (Young *et al.* 1997; Field *et al.* 2000; Halpin *et al.* 2000). Our data demonstrate that HeV risk is high in all reproducing females, including lactating and near-term pregnant animals (animals that have not aborted their foetus), suggesting that increased numbers of infected animals, as opposed to simply abortion events, could drive the higher force of infection to spillover hosts during the reproductive season.

In the autumn 2006 birthing period, no females were pregnant or lactating. Coincident with this period was a shortage of blossom and nectar, presumably due to high rainfall and a prolonged wet season. Low body weights, poor body conditions and observations of extremely abnormal diurnal feeding indicate that the population was nutritionally stressed, which probably accounts for the non-reproductive state of adult females at this time. Despite the lack of reproduction, all demographic groups exhibited approximately 80% seroprevalence, 14–42 times higher than the odds of seropositivity in any other season after adjusting for sex, reproductive status and age (table 3). The elevated seroprevalence in a nutritionally stressed population could be explained by increased viral




Figure 4. Relationship between seroprevalence and body condition (weight/forearm ratio). Points represent mean seroprevalence per field season as a function of mean body weight/forearm ratio for adult non-pregnant females (squares and solid line) and adult males (diamonds and dashed line). Dividing body weight by forearm measurement accounts for body size and therefore gives a crude measurement of body condition. 1, spring 2004; 2, autumn 2005; 3, winter 2005; 4, spring 2005; 5, autumn 2006.

susceptibility due to substantial energetic costs of immune responses (Gulland 1992; Sheldon & Verhulst 1996; Nelson 2004). Furthermore, nutritional and environmental stresses can change behaviour, for example causing animals to crowd around restricted food resources or to share food with other species. Our data (figure 4) demonstrate that body condition and seroprevalence were related in all field seasons: seroprevalence increased when body condition decreased, with the exception of males during the two mating seasons.

We hypothesized that mating would be associated with increased risk of viral transmission, because it is a time of intense interaction (Markus 2002) and the immunosuppressive effects of testosterone have been shown to increase disease risk (Zuk & McKean 1996). In the spring 2004 mating season, there was a high population seroprevalence (figure 3*b*), particularly in adult males. However, during the spring 2005 mating season, we observed low seroprevalence in all animals, yet mating was observed, and age distributions and population size were comparable with those of spring 2004. A plausible explanation for our results from spring 2005 is that HeV infection was not present in the population at that time. This interpretation is supported by modelling efforts of HeV infection dynamics in the other flying fox species, which suggest that HeV does not persist endemically within spatially discrete local populations, but persists globally via metapopulation dynamics (Plowright 2007).

Although the absence of infection provides a plausible explanation for the low HeV seroprevalence in LRFF in spring 2005, it is an improbable explanation for the marked decline in seroprevalence we observed between autumn 2005 and winter/spring 2005 (figure 3*b*). It may be that, unlike detectable immunity in other flying fox species, detectable immunity in LRFF wanes over short time scales. IgG to HeV in *P. conspicillatus* and *P. poliocephalus* is detectable for years-to-a-lifetime (Field




Figure 5. Variance in HeV seroprevalence over five field seasons in LRFF compared with variance in HeV seroprevalence over five field seasons in a Northern Queensland population of spectacled flying foxes (A. C. Breed 2005–2006, unpublished data).

2005; H. Field, unpublished data), resulting in relatively stable seroprevalence (figure 5) over time (Field 2005).

Waning immunity would significantly enhance the persistence of HeV in LRFF, conferring on them a critical role in the maintenance of HeV across *Pteropus* spp. Given that LRFF resolve as the most basal lineage of *Pteropus* bats (O'Brien 2005), and are genetically distant (Bastian *et al.* 2002; O'Brien 2005), and ecologically and morphologically distinct (Hall & Richards 2000; Birt 2004), from other Australian flying fox species, waning immunity may reflect a more mature host–pathogen relationship in this species.

In our study, flying fox campsites moved in relation to ephemeral patches of blossom and nectar; therefore, we sampled in multiple locations. In our analysis, we assumed that individuals sampled at different locations were part of a single population and that the patterns observed were not due to sampling of different populations. We based this assumption on: (i) almost identical seroprevalence among locations within seasons, suggesting regional synchrony of viral dynamics, (ii) satellite and radio telemetry studies which indicated considerable movement of individuals among roost sites within the study area (R. K. Plowright, C. Smith, C. Palmer, A. Divljan & H. Field 2005, unpublished data; C. Palmer, C. Smith, A. Divljan & R. K. Plowright 2005, unpublished data), and (iii) the rapid decline in seroprevalence observed between autumn and winter 2005, which was probably not due to sampling of different populations, as we sampled at the same location consecutively during those two seasons. Furthermore, we are confident that at least the lactating females in autumn and winter 2005 represent a single population, because lactating females typically do not move during pup dependency (Birt 2004) and we observed the same pattern of declining seroprevalence in these animals. We therefore conclude that disparate local populations may actually constitute one regional population for viral dynamics, that sampling different populations did not confound our results, and that the overall rapid decline in

seroprevalence observed between seasons is not due to sampling of different populations.

In summary, our research identified a pattern of horizontal transmission, probably through urine, faeces or saliva, with increasing age significantly increasing risk of infection through direct contact. Excreta- or saliva-contaminated pastures, or discarded fruit, are probable routes of viral transmission to horses, and our data suggest that risk of transmission increases during flying fox reproductive periods and in times of nutritional stress. Our conclusion that LRFF immune dynamics seem to differ from dynamics in the three other Australian *Pteropus* species may be critical for understanding viral dynamics in all *Pteropus* spp.

Increased infection risk in reproductively and nutritionally stressed animals may have important ramifications for the dynamics of henipaviruses in *Pteropus* species. First, it suggests that there is probably an element of seasonality to infection dynamics that may be useful in spillover prevention and control. Second, anthropogenic habitat loss, habitat alteration, roost disturbance, urbanization and persecution by hunters (Fujita & Tuttle 1991; Mickleburg *et al.* 1992; Duncan *et al.* 1999), all of which are known to alter food availability and cause nutritional stress in *Pteropus* populations (Hall & Richards 2000; P. Eby 1998–2007, unpublished data), may drive HeV spillover events. Finally, nectar and fruit availability are dependent on climatic conditions (Law *et al.* 2000), and thus flying foxes—particularly the highly nectar-dependent LRFF—may be greatly affected by climate change. Our data suggest that such environmental stresses lead directly to changes in henipavirus infection and excretion dynamics. Future work to test the hypothesis that nutritional stress, driven by climate change and other anthropogenic alterations to the environment, is an important driver of disease emergence for HeV, and perhaps other emerging bat viruses, is urgently needed.

Charles Darwin University Animal Ethics Committee gave permission for work on *P. scapulatus* (project reference: A04033). Permits were granted by the Parks and Wildlife Commission of the Northern Territory (permit number: 18597).

We thank the Northern Territory Government's Department of Natural Resources, Environment and the Arts, particularly Katherine region staff and John Woinarski from Biodiversity Conservation Division, Darwin, for their great support during the project. We also thank Greer Meehan, Rhys Fogarty, John Burke, Justin Perry, Rhonda Scocinarro, Tracey Blackney, Sam Veloz, Liz Chamberlin, Adam Porter, Damian Milne, Dave Fuller, Carol de Jong, Kerryn Parry-Jones, Marion Cook, Peter Cook, Chris Kinard, David Hooper, Andy Peckham, Amanda McLaughlin, Sam Wushusen and Katharine Bossart for their assistance with the field and laboratory work; Les Hall, Patrina Birt, Peggy Eby, Kim Halpin, Juliet Pulliam, Patrick Foley, Phil Kass, Andrew Breed, Alex Hyatt, Deborah Middleton, Andy Dobson, Jon Epstein, Lin-Fa Wang, Bryan Eaton, Andrew Cunningham and Kevin Olival for their insightful conversations about LRFF and HeV; Rosie Woodroffe, Bruno Chomel, Leslie Bienen, Paul Cross, John Winnie Jr and Sara Gregory for their comments on the manuscript.

This work was supported in part by an NIH/NSF 'Ecology of Infectious Diseases' award from the John E. Fogarty International Center R01-TW05869, by core funding to the Consortium for Conservation Medicine from the V. Kann Rasmussen Foundation and is published in collaboration with

the Australian Biosecurity Cooperative Research Center for Emerging Infectious Diseases (AB-CRC). R.K.P. was supported by the V. Kann Rasmussen Foundation, the Australian-American Fulbright Commission, the Foundation for Young Australians and grants from the UC Davis Graduate Group in Ecology.

REFERENCES

- Alexander, J. & Stimson, W. H. 1988 Sex-hormones and the course of parasitic infection. *Parasitol. Today* **4**, 189–193. (doi:10.1016/0169-4758(88)90077-4)
- Bastian Jr, S. T., Tanaka, K., Anunciado, R. V. P., Natural, N. G., Sumalde, A. C. & Namikawa, T. 2002 Evolutionary relationships of flying foxes (Genus *Pteropus*) in The Philippines inferred from DNA sequences of cytochrome *b* gene. *Biochem. Genet.* **40**, 101–116. (doi:10.1023/A:1015161305843)
- Birt, P. 2004 Mutualistic interactions between the nectar-feeding little red flying fox *Pteropus scapulatus* (Chiroptera: Pteropodidae) and flowering eucalypts (Myrtaceae): habitat utilisation and pollination. PhD thesis, University of Queensland, Brisbane, Australia.
- Bossart, K. N., McEachern, J. A., Hickey, A. C., Choudhry, V., Dimitrov, D. S., Eaton, B. T. & Wang, L. F. 2007 Neutralization assays for differential henipavirus serology using Bio-Plex protein array systems. *J. Virol. Methods* **142**, 29–40. (doi:10.1016/j.jviromet.2007.01.003)
- Cattadori, I. M., Boag, B., Bjornstad, O. N., Cornell, S. J. & Hudson, P. J. 2005 Peak shift and epidemiology in a seasonal host-nematode system. *Proc. R. Soc. B* **272**, 1163–1169. (doi:10.1098/rspb.2004.3050)
- Chua, K. B. *et al.* 2007 A previously unknown reovirus of bat origin is associated with an acute respiratory disease in humans. *Proc. Natl Acad. Sci. USA* **104**, 11 424–11 429. (doi:10.1073/pnas.0701372104)
- Cohen, M. L. 2000 Changing patterns of infectious disease. *Nature* **406**, 762–767. (doi:10.1038/35021206)
- Daniels, P., Ksiazek, T. & Eaton, B. T. 2001 Laboratory diagnosis of Nipah and Hendra virus infections. *Microbes Infect.* **3**, 289–295. (doi:10.1016/S1286-4579(01)01382-X)
- Daszak, P., Cunningham, A. A. & Hyatt, A. D. 2000 Emerging infectious diseases of wildlife—threats to biodiversity and human health. *Science* **287**, 443–449. (doi:10.1126/science.287.5452.443)
- Daszak, P. *et al.* 2006 The emergence of Nipah and Hendra virus: pathogen dynamics across a wildlife–livestock–human continuum. In *Disease ecology: community structure and pathogen dynamics* (eds S. Collinge *et al.*), pp. 186–201. Oxford, UK: Oxford University Press.
- Divljan, A., Parry-Jones, K. & Wardle, G. M. 2006 Age determination in the grey-headed flying fox. *J. Wildl. Manage.* **70**, 607–611. (doi:10.2193/0022-541X(2006)70[607:ADITGF]2.0.CO;2)
- Dobson, A. & Foufopoulos, J. 2001 Emerging infectious pathogens of wildlife. *Phil. Trans. R. Soc. B* **356**, 1001–1012. (doi:10.1098/rstb.2000.0758)
- Duncan, A., Baker, G. B. & Montgomery, N. 1999 *The action plan for Australian bats*. Canberra, Australia: Environment Australia.
- Eaton, B. T., Broder, C. C., Middleton, D. & Wang, L. F. 2006 Hendra and Nipah viruses: different and dangerous. *Nat. Rev. Microbiol.* **4**, 23–35. (doi:10.1038/nrmicro1323)
- Epstein, J. H., Field, H. E., Luby, S., Pulliam, J. R. & Daszak, P. 2006 Nipah virus: impact, origins, and causes of emergence. *Curr. Infect. Dis. Rep.* **8**, 59–65. (doi:10.1007/s11908-006-0036-2)

- Festa-Bianchet, M. 1989 Individual differences, parasites, and the costs of reproduction for bighorn ewes (*Ovis canadensis*). *J. Anim. Ecol.* **58**, 785–795. (doi:10.2307/5124)
- Field, H. 2005 The ecology of Hendra virus and Australian bat lyssavirus. PhD thesis, The University of Queensland, Brisbane.
- Field, H., Barrat, P. C., Hughes, R. J., Shield, J. & Sullivan, N. D. 2000 A fatal case of Hendra virus infection in a horse in north Queensland—clinical and epidemiological features. *Aust. Vet. J.* **78**, 279–280.
- Field, H., Young, P., Yob, J. M., Mills, J., Hall, L. & Mackenzie, J. 2001 The natural history of Hendra and Nipah viruses. *Microbes Infect.* **3**, 307–314. (doi:10.1016/S1286-4579(01)01384-3)
- Field, H. E., Breed, A. C., Shield, J., Hedlefs, R. M., Pittard, K., Pott, B. & Summers, P. M. 2007 Epidemiological perspectives on Hendra virus infection in horses and flying foxes. *Aust. Vet. J.* **85**, 268–270. (doi:10.1111/j.1751-0813.2007.00170.x)
- Fujita, M. S. & Tuttle, M. D. 1991 Flying foxes (Chiroptera: Pteropodidae): threatened animals of key ecological and economic importance. *Conserv. Biol.* **5**, 455–463. (doi:10.1111/j.1523-1739.1991.tb00352.x)
- Gulland, F. M. 1992 The role of nematode parasites in Soay sheep (*Ovis aries* L.) mortality during a population crash. *Parasitology* **105**(Pt 3), 493–503.
- Hall, L. & Richards, G. 2000 *Flying foxes: fruit and blossom bats of Australia*. Australian natural history series. Sydney, Australia: University of New South Wales Press Ltd.
- Halpin, K., Young, P. L., Field, H. E. & Mackenzie, J. S. 2000 Isolation of Hendra virus from *Pteropid* bats: a natural reservoir of Hendra virus. *J. Gen. Virol.* **81**, 1927–1932.
- Hanna, J. N., McBride, W. J., Brookes, D. L., Shield, J., Taylor, C. T., Smith, I. L., Craig, S. B. & Smith, G. A. 2006 Hendra virus infection in a veterinarian. *Med. J. Aust.* **185**, 562–564.
- Jonsson, N. N., Johnston, S. D., Field, H., de Jong, C. & Smith, C. 2004 Field anaesthesia of three Australian species of flying fox. *Vet. Rec.* **154**, 664.
- Law, B., Mackowski, C., Schoer, L. & Tweedie, T. 2000 Flowering phenology of myrtaceous trees and their relation to climatic, environmental and disturbance variables in northern New South Wales. *Austral Ecol.* **25**, 160–178. (doi:10.1046/j.1442-9993.2000.01009.x)
- Lloyd, S. S. 1983 Immunosuppression during pregnancy and lactation. *Ir. Vet. J.* **37**, 64–70.
- Mackenzie, J. S. *et al.* 2001 Emerging viral diseases of Southeast Asia and the Western Pacific. *Emerg. Infect. Dis.* **7**, 497–504.
- Mackenzie, J. S., Field, H. E. & Guyatt, K. J. 2003 Managing emerging diseases borne by fruit bats (flying foxes), with particular reference to henipaviruses and Australian bat lyssavirus. *J. Appl. Microbiol.* **94**, 59s–69s. (doi:10.1046/j.1365-2672.94.s1.7.x)
- Markus, N. 2002 Behaviour of the black flying fox *Pteropus alecto*: 2. Territoriality and courtship. *Acta Chiropt.* **4**, 153–166.
- Mickleburg, S., Hutson, A. & Racey, P. 1992 *Old world fruit bats: an action plan for their conservation*. Gland, Switzerland: International Union for the Conservation of Nature and Natural Resources (IUCN).
- Murray, K. *et al.* 1995 A morbillivirus that caused fatal disease in horses and humans. *Science* **268**, 94–97. (doi:10.1126/science.7701348)
- Nelson, R. J. 2004 Seasonal immune function and sickness responses. *Trends Immunol.* **25**, 187–192. (doi:10.1016/j.it.2004.02.001)
- O'Brien, J. 2005 Phylogeography and conservation genetics of the fruit bat genus *Pteropus* (Megachiroptera) in the western Indian Ocean. PhD thesis, National University of Ireland, Dublin.
- Palmer, C. & Woinarski, J. C. Z. 1999 Seasonal roosts and foraging movements of the black flying fox (*Pteropus alecto*) in the Northern Territory: resource tracking in a landscape mosaic. *Wildl. Res.* **26**, 823–838. (doi:10.1071/WR97106)
- Palmer, C., Price, O. & Bach, C. 2000 Foraging ecology of the black flying fox (*Pteropus alecto*) in the seasonal tropics of the Northern Territory Australia. *Wildl. Res.* **27**, 169–178. (doi:10.1071/WR97126)
- Plowright, R. K. 2007 The ecology and epidemiology of Hendra virus in flying foxes (*Pteropus* spp.). PhD thesis, University of California, Davis.
- Plowright, R. K., Sokolow, S. H., Gorman, M. E., Daszak, P. & Foley, J. E. 2008 Causal inference in disease ecology: investigating ecological drivers of disease emergence. *Front. Ecol. Environ.* **6**, (doi:10.1890/070086)
- R Development Core Team 2006 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Sheldon, B. C. & Verhulst, S. 1996 Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends Ecol. Evol.* **11**, 317–321. (doi:10.1016/0169-5347(96)10039-2)
- Smolinski, M. S., Hamburg, M. A. & Lederberg, J. 2003 *Microbial threats to health: emergence, detection, and response*. Washington, DC: The National Academies Press.
- Trevelo, R. T., Barr, M. C. & Robinson, R. A. 2005 Important emerging bacterial zoonotic infections affecting the immunocompromised. *Vet. Res.* **36**, 493–506. (doi:10.1051/vetres:2005011)
- Williamson, M. M., Hooper, P. T., Selleck, P. W., Gleeson, L. J., Daniels, P. W., Westbury, H. A. & Murray, P. K. 1998 Transmission studies of Hendra virus (equine morbillivirus) in fruit bats, horses and cats. *Aust. Vet. J.* **76**, 813–818.
- Williamson, M. M., Hooper, P. T., Selleck, P. W., Westbury, H. A. & Slocumbe, R. F. 1999 Experimental Hendra virus infection in pregnant guinea-pigs and fruit bats (*Pteropus poliocephalus*). *J. Comp. Pathol.* **122**, 201–207. (doi:10.1053/jcpa.1999.0364)
- Young, P., Halpin, K., Field, H. & Mackenzie, J. 1997 Finding the wildlife reservoir of equine morbillivirus. In *Recent advances in microbiology*, vol. 5 (ed. V. Asche), pp. 1–12. Melbourne, Australia: Australian Society of Microbiology, Inc.
- Zuk, M. & McKean, K. A. 1996 Sex differences in parasite infections: patterns and processes. *Int. J. Parasitol.* **26**, 1009–1023. (doi:10.1016/S0020-7519(96)00086-0)