

Correction



Cite this article: Ruta M, Angielczyk KD, Fröbisch J, Benton MJ. 2013 Decoupling of morphological disparity and taxic diversity during the adaptive radiation of anomodont therapsids. *Proc R Soc B* 280: 20132414. <http://dx.doi.org/10.1098/rspb.2013.2414>

Decoupling of morphological disparity and taxic diversity during the adaptive radiation of anomodont therapsids

Marcello Ruta, Kenneth D. Angielczyk, Jörg Fröbisch and Michael J. Benton

Proc. R. Soc. B 280, 20131071 (7 October 2013; Published online 14 August 2013)
(doi:10.1098/rspb.2013.1071)

Figure 3*h* in the original paper contains an inaccuracy. The bottom right panel (3*h*) is identical to panel 3*g*. The correct panel 3*h*, describing the plot of the rarefied distance from founder, is reproduced here together with all other panels of figure 3. The caption of that figure is slightly altered.




Figure 3. (a–d) Comparisons between anomodont disparity (rarefied median values, grey circles) and diversity (white squares) through time intervals t1–t8; the disparity values are for the sum of ranges (a,c) and the sum of variances (b,d); the error bars around the diversity values are calculated as $\pm \sqrt{N}$, where N is the number of taxa in any given interval; (a,b) comparisons based on the number of taxa present in the phylogeny; (c,d) comparisons based on the total number of known anomodont taxa. (e–h) Distance from the founder (white rhombs), expressed as the average generalized Euclidean distance of taxa from *Biseridens* ('founder' taxon); (e,f) average distance of taxa binned by time intervals; (g,h) average distance of major groups. Vertical grey bars mark the Permian–Triassic boundary.