



Research

Cite this article: Pamerter ME, Dzal YA, Milsom WK. 2015 Adenosine receptors mediate the hypoxic ventilatory response but not the hypoxic metabolic response in the naked mole rat during acute hypoxia. *Proc. R. Soc. B* **282**: 20141722.

<http://dx.doi.org/10.1098/rspb.2014.1722>

Received: 10 July 2014

Accepted: 20 November 2014

Subject Areas:

physiology

Keywords:

hypoxic ventilatory response, hypoxic metabolic response, plethysmography, respirometry, thermal response

Author for correspondence:

Matthew E. Pamerter

e-mail: pamerter@zoology.ubc.ca

[†]These authors contributed equally to this study.

Adenosine receptors mediate the hypoxic ventilatory response but not the hypoxic metabolic response in the naked mole rat during acute hypoxia

Matthew E. Pamerter[†], Yvonne A. Dzal[†] and William K. Milsom

Department of Zoology, University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z4

Naked mole rats are the most hypoxia-tolerant mammals identified; however, the mechanisms underlying this tolerance are poorly understood. Using whole-animal plethysmography and open-flow respirometry, we examined the hypoxic metabolic response (HMR), hypoxic ventilatory response (HVR) and hypoxic thermal response in awake, freely behaving naked mole rats exposed to 7% O₂ for 1 h. Metabolic rate and ventilation each reversibly decreased 70% in hypoxia (from 39.6 ± 2.9 to 12.1 ± 0.3 ml O₂ min⁻¹ kg⁻¹, and 1412 ± 244 to 417 ± 62 ml min⁻¹ kg⁻¹, respectively; $p < 0.05$), whereas body temperature was unchanged and animals remained awake and active. Subcutaneous injection of the general adenosine receptor antagonist aminophylline (AMP; 100 mg kg⁻¹, in saline), but not control saline injections, prevented the HVR but had no effect on the HMR. As a result, AMP-treated naked mole rats exhibited extreme hyperventilation in hypoxia. These animals were also less tolerant to hypoxia, and in some cases hypoxia was lethal following AMP injection. We conclude that in naked mole rats (i) hypoxia tolerance is partially dependent on profound hypoxic metabolic and ventilatory responses, which are equal in magnitude but occur independently of thermal changes in hypoxia, and (ii) adenosine receptors mediate the HVR but not the HMR.

1. Introduction

When O₂ supply fails to meet O₂ demand, cellular energy balance is disrupted, which can ultimately lead to cell death, organ failure and mortality. Therefore, matching O₂ supply to O₂ demand is critical to tolerating hypoxia [1]. Hypoxic environments are common on Earth, and exposure to periods of intermittent or prolonged hypoxia in the daily activities or annual life cycles of various animals has driven the evolution of diverse cellular, physiological and behavioural adaptations to low O₂ stress [2]. These adaptations typically serve to increase O₂ supply to cells and tissues, and/or to decrease cellular energy demand (i.e. metabolic rate depression) [2,3]. For example, when most mammals are exposed to acute hypoxia, their immediate compensatory response is to increase their ventilation [4], which is an energetically expensive means of increasing the delivery of O₂ to cells. Conversely, some species decrease breathing in hypoxia, although this is not common in mammals. Such hypoxia-mediated changes in breathing are termed the hypoxic ventilatory response (HVR), which encompasses a suite of time domains depending on the duration of hypoxic exposure [4]. Within a few moments of the onset of the HVR, most mammals also exhibit a decrease in metabolic demand, termed the hypoxic metabolic response (HMR), which in turn decreases systemic O₂ requirements. As metabolic demand decreases, ventilation (\dot{V}_E , expiratory minute ventilation) tends to decrease as well; however, in all mammals studied to date, the decrease in metabolism is greater than the decrease in \dot{V}_E . Therefore, most mammals tend to hyperventilate in hypoxia because the HVR and HMR combine to affect a net increase in the animals' air convection requirement (ACR; i.e. the ratio of inspired O₂ to metabolic rate).

The magnitude of the HVR and HMR are key determinants of the duration and depth of hypoxia that an organism can tolerate. Certain air-breathing vertebrates, such as arctic ground squirrels (*Urocyon parryi*); freshwater turtles (*Chrysemys picta bellii*) and crucian carp (*Carassius carassius*), can tolerate hypoxia for weeks to months without apparent detriment [2]. Typically, these animals are most hypoxia-tolerant while over-wintering. During this period, they exhibit metabolic rate depression mediated in part by a profound decrease in body temperature (T_b) and entry into a torpor-like state [5]. These responses drastically reduce systemic energy demands and extend survival time in hypoxia. The mechanisms that mediate these changes are poorly understood, but recent studies have demonstrated that in small mammals, adenosine receptors are key in initiating and maintaining (i) the HMR, (ii) decreases in T_b and (iii) the entry into torpor. In support of this, adenosine receptor agonism decreases metabolism and T_b , and induces entry into a torpor-like state in rats and arctic ground squirrels [6,7], whereas adenosine receptor antagonism prevents or reverses these changes in hibernating arctic ground squirrels [7,8], and during intra-day torpor or fasting-induced torpor in mice [9–11] and rats [6].

Naked mole rats (*Heterocephalus glaber*) experience lifelong hypoxia while living in large subterranean tunnel systems approximately 1–2 m underground in eastern Africa [12]. Naked mole rat burrows are poorly ventilated due to limited air diffusion through densely packed soil, which, in combination with group respiration-mediated O_2 depletion, creates a challenging hypoxic environment [12,13]. Indeed, naked mole rats and their relatives experience burrow O_2 levels as low as 6.2% without detriment [12–14] and tolerate hypoxia lower than 3% O_2 for several hours in a laboratory setting [15]. Based on these observations, naked mole rats are considered to be the most hypoxia-tolerant mammal. Owing to the constancy of their exposure to environmental hypoxia and also the relatively high ambient temperature (T_a) in their burrows (approx. 30–34°C), naked mole rats cannot easily use behavioural and thermal means to alter their metabolic rate. Nonetheless, studies to date have described a handful of adaptations in naked mole rats that are likely to contribute to their tolerance of chronic hypoxia. These include the expression of a haemoglobin isomer with higher affinity for O_2 relative to other rodents [16], a metabolic rate that is two-thirds of that predicted from allometry [17], and blunted neuronal Ca^{2+} influx during hypoxia [18–20], which is known to reduce excitotoxic cell death due to hypoxia or ischaemia [21]. However, even the most basic system-level responses, including ventilatory and metabolic responses to hypoxia, remain to be properly characterized in this species. The aims of this study were to examine (i) the steady-state metabolic, ventilatory and thermal responses of naked mole rats to acute hypoxia, and (ii) the role of adenosinergic signalling in mediating these responses.

2. Material and methods

(a) Animals

Naked mole rats were generous gifts from the University of Montana and the University of Texas at San Antonio. Naked mole rats were group-housed in interconnected multi-cage systems and held at 28°C in 50% humidity under a 12 L:12 D cycle. Animals were fed fresh tubers, vegetables, fruit and Pronutro cereal supplement ad libitum. Animals were not fasted prior to the experimental trial.

(b) Experimental design

Fourteen male and female naked mole rats weighing 45.1 ± 1.8 g were individually placed, unrestrained, into an experimental chamber flushed with 21% O_2 , which was set inside a controlled environmental chamber to maintain T_a at approximately 28°C. Once placed within the experimental apparatus, naked mole rats were given at least 30 min to acclimate, followed by 60 min of baseline recordings of \dot{V}_E and metabolism. This period was necessary to stabilize the chamber temperature and to determine steady-state metabolism and \dot{V}_E . After baseline recordings, the naked mole rat was quickly removed from the chamber and its T_b was measured rectally using a thermocouple. Naked mole rats were then randomly divided into two groups and treated with subcutaneous 500 μ l injections of either (i) saline ($n = 7$) or (ii) the general adenosine receptor antagonist aminophylline (AMP) dissolved in saline (100 mg kg^{-1} ; $n = 7$). AMP was purchased from Sigma-Aldrich (St Louis, MO, USA). In the body, AMP is converted to theophylline with approximately 80% efficacy [22]. The pharmacokinetics of intravenous, intraperitoneal or subcutaneous AMP administration best fitted the two-compartment model, with rapid distribution and slow elimination. Titres of plasma theophylline peak within 8 min of injection and the normal elimination half-life of AMP ranges from 8 to 14 h, while the bioavailability of AMP is 80–95% in humans [23], dogs [24], cats [25], pigs [26] and chickens [25]. Although the pharmacokinetics and bioavailability of AMP have not been previously assessed in naked mole rats, these are likely to be similar to published values in other species. Furthermore, the pharmacological dose of AMP used in this study is consistent with that used to inhibit metabolic depression in studies conducted in small rodents in other laboratories (e.g. [11]). We observed no negative effects of AMP administration. Specifically, animals did not appear stressed, their behaviour was not altered, and their metabolic and ventilatory rates were unaffected. Immediately following AMP administration, animals were returned to the recording chamber, and ventilatory and metabolic measurements were collected in each gas mixture (21% O_2 , 7% O_2 and recovery in 21% O_2), under poikilothermic conditions. The effects of AMP were determined by comparing data obtained from control animals (saline injection) relative to data collected from animals treated with AMP. Animals were held in each condition until metabolism and \dot{V}_E reached steady state (approx. 1 h), and T_b was measured following each exposure. The experimental trial took a total of approximately 4 h for each individual.

(c) Plethysmography and respirometry

\dot{V}_E was measured using the barometric pressure method of plethysmography modified for continuous flow [27]. On the day of experimentation, individuals were placed into the experimental apparatus which consisted of two rectangular 450 ml Plexiglas chambers, one being the animal chamber and the other the reference chamber. The temperature of the animal chamber was recorded continuously throughout the experiment using an iButton (Maxim Integrated, Chandler, CA). Calibrated rotameters were used to supply naked mole rats with an inflowing gas mixture set at a flow rate of 110 $ml\ min^{-1}$. The total airflow ensured that neither O_2 nor CO_2 were altered by more than 1.5% by the animal's metabolism. Fractional O_2 and CO_2 composition of inspired and expired gas were monitored using an O_2 and CO_2 analyser (Raytech, North Vancouver, BC, Canada). The gas analyser was calibrated for O_2 and CO_2 before each trial with a premixed gas (21% O_2 ; 1.55% CO_2 balanced with N_2). O_2 consumption was calculated from the product of the constant airflow through the chamber and the difference between the inflow and outflow in the fractional concentration of O_2 . CO_2 production was calculated from the product of the constant airflow through the chamber and the difference between the outflow and inflow in the fractional concentration of CO_2 .

The respiratory exchange ratio (RER) was calculated as the quotient of CO₂ production and O₂ consumption. The ACR was calculated as the quotient of \dot{V}_E and O₂ consumption. Inspiration humidifies and warms inspired air, resulting in pressure fluctuations that were compared with the pressure of the reference chamber and monitored with a differential pressure transducer (DP103-18, Validyne, Northridge, CA, USA) connected between the two chambers, amplified and electronically recorded by a computer. Respiratory frequency (f_R) was calculated directly from the ventilation-induced pressure oscillations. Expiratory tidal volume (V_T) was determined by integrating negative periods of ventilatory flow and then calculated using the equation of Drorbaugh & Fenn [28] modified for flow-through plethysmography by Jacky [27]. To determine V_T , the system was calibrated dynamically with the animal present in the chamber, as described previously [29]. Briefly, prior to each experiment known volumes of air (0.2, 0.5 and 1.0 ml) were injected into the chamber at a rate similar to the naked mole rats' expiratory time to produce pressure deflections at least 10 times as great as that of the animal's breathing. \dot{V}_E was measured with the system open and oscillations in recorded pressure were considered to be proportional to V_T . The advantage of an open-flow system is that the animal chamber is flushed constantly with air so that O₂ depletion and CO₂ accumulation are avoided, and metabolism and \dot{V}_E could be continuously and simultaneously recorded. \dot{V}_E was calculated from the product of f_R and V_T .

(d) Data collection and analysis

Ventilatory, metabolic and thermoregulatory variables were recorded on WINDAQ acquisition software (Dataq Instruments, Akron, OH, USA). All data acquired in WINDAQ were analysed in POWERLAB. Average values were calculated for each variable during the last 5 min of steady state of each experimental treatment. For \dot{V}_E , V_T and f_R , a series of at least 10 sets of 10 consecutive breaths were selected for analysis. Breathes were chosen from periods where the animal was awake but not active (i.e. not actively digging or exploring but also not asleep, as determined by visual examination).

Statistical analysis was performed using commercial software (SPSS v. 15.0; SPSS Inc., Chicago, IL, USA). For all experiments, individual n values correspond to a single animal treated with either saline or AMP dissolved in saline as described above, before and after exposure to 21% O₂ and then acute hypoxia (7% O₂). Values are presented as mean \pm s.e.m. $p < 0.05$ was considered to achieve statistical significance. All data were normally distributed with equal variance ($p > 0.05$). Significance was evaluated using two-way ANOVA to test for significant interactions between the two independent variables: (i) the acute inspired O₂ level (7% versus 21%) and (ii) the treatment (saline versus AMP injection). Bonferroni post hoc multiple-comparison tests were run on each of the dependent variables to compare the single point means of interest. The dependent variables analysed were \dot{V}_E , f_R , V_T , metabolic rate, RER, the ACR and T_b .

3. Results

(a) Naked mole rats exhibit robust metabolic depression with sustained T_b in acute hypoxia

Steady-state metabolism, \dot{V}_E , and T_b were measured in naked mole rats breathing normoxic gas mixtures before and after subcutaneous saline or AMP injection. Neither saline nor AMP altered any of these variables in animals breathing normoxic gas mixtures ($p > 0.05$; $n = 7$ each; data not shown). Thus, for the purposes of data presentation and statistical analysis we used post-injection normoxic measurements as the control in

all datasets. Relative to control levels in normoxia, saline-treated naked mole rats exposed to 7% O₂ exhibited a 70% decrease in metabolic rate (from 39.6 ± 2.9 to 12.1 ± 0.3 ml O min⁻¹ kg⁻¹, and 31.3 ± 2.7 to 8.7 ± 1.1 ml CO₂ min⁻¹ kg⁻¹, for O₂ consumption and CO₂ production, respectively; figure 1*a,b*), which was reversed following 30 min of recovery in normoxia (21% O₂; to 45.4 ± 4.6 ml O₂ min⁻¹ kg⁻¹ and 23.8 ± 3.4 ml CO₂ min⁻¹ kg⁻¹). Similarly, AMP-treated naked mole rats breathing 7% O₂ also exhibited metabolic rate depression of a similar magnitude (from 38.6 ± 2.3 to 15.1 ± 3.8 ml O₂ min⁻¹ kg⁻¹, and 28.3 ± 1.6 to 17.2 ± 2.5 ml CO₂ min⁻¹ kg⁻¹, for O₂ consumption and CO₂ production, respectively; figure 1*a,b*), although the hypoxic decrease in CO₂ production in AMP-treated naked mole rats was significantly less than that observed in saline-treated animals (figure 1*b*). This difference may have been due to a secondary effect of AMP on the balance between the catabolism of lipids and carbohydrates as fuel substrates in hypoxic naked mole rats, although this requires further investigation [30,31]. These changes were also reversed following 30 min of recovery in normoxia (to 41.1 ± 3.3 ml O₂ min⁻¹ kg⁻¹ and 22.5 ± 0.8 ml CO₂ min⁻¹ kg⁻¹). T_b was not different between treatments or inspired O₂ levels in any group (figure 1*c*). Visual observations revealed that naked mole rats remained alert and active throughout the experiment in all treatment groups.

(b) Naked mole rats exhibit ventilatory depression in acute hypoxia mediated by adenosine receptors

When breathing normoxic gas mixtures, \dot{V}_E was 1412 ± 224 ml min⁻¹ kg⁻¹ in saline-treated animals (figure 2*a*). Following the switch to 7% O₂, \dot{V}_E decreased by 70% to 417 ± 62 ml min⁻¹ kg⁻¹. This value reversed to 1071 ± 259 ml min⁻¹ kg⁻¹ after 30 min of recovery in normoxia, which was not significantly different from control values. The change in \dot{V}_E in 7% O₂ was due to the combination of a 50% decrease in f_R (from 120.1 ± 15.8 to 59.9 ± 4.5 breaths min⁻¹; figure 2*b*) and a 47% decrease in V_T (from 13.6 ± 1.7 to 7.2 ± 1.3 ml kg⁻¹; figure 2*c*), both of which were reversed by reoxygenation. In AMP-treated animals, \dot{V}_E in normoxia was not different from saline-treated animals (1319 ± 260 ml min⁻¹ kg⁻¹; figure 2*a*); however, the hypoxia-mediated decrease in \dot{V}_E did not occur and \dot{V}_E in AMP-treated animals breathing 7% O₂ was 1344 ± 205 ml min⁻¹ kg⁻¹. In these naked mole rats, the hypoxia-mediated decrease in f_R was not observed and f_R tended to increase throughout the hypoxic exposure and recovery, although this trend did not reach significance (figure 2*b*). Conversely, the hypoxia-mediated decrease in V_T was only partially prevented by AMP treatment, and the observed V_T in these animals was statistically different from both pre-hypoxia control values in AMP-treated animals and also from hypoxic values in saline-treated animals breathing 7% O₂ (figure 2*c*). In addition, the hypoxic decrease in V_T did not reverse upon reoxygenation in AMP-treated animals.

(c) Hypoxia does not alter the air convection requirement in naked mole rats

The ACR for both O₂ consumption and CO₂ production indicated that saline-treated naked mole rats did not exhibit a change in breathing relative to metabolism when exposed to 7% O₂ (ACR_{O₂}: 35.6 ± 4.6 in normoxia versus 34.5 ± 6.8 in hypoxia; ACR_{CO₂}: 45.1 ± 3.2 in normoxia versus 47.9 ± 4.9 in hypoxia; figure 3*a,b*). This was expected, as the




Figure 1. Metabolism is markedly reduced during acute hypoxia independently of ADO receptor activation. (a,b) Average metabolic rate in naked mole rats breathing normoxic or acute hypoxic (7% O₂) gas mixtures and treated with subcutaneous saline (white bars) or AMP (black bars) injection. (c) Average body temperature from the animals treated in (a,b). Data are mean \pm s.e.m. from $n = 7$ naked mole rats per group. Asterisks (*) indicate significant differences between normoxia and acute hypoxia values; daggers (†) indicate significant differences between AMP- and saline-treated animals ($p < 0.05$).




Figure 2. Ventilation is markedly reduced during acute hypoxia in an ADO receptor-dependent manner. Effects of acute hypoxia (7% O₂), saline (white circles, dashed lines) and AMP (black squares, solid lines) on (a) total minute ventilation (V_E), (b) breathing frequency (f_R) and (c) tidal volume (V_T) in naked mole rats. Data are mean \pm s.e.m. from $n = 7$ naked mole rats per group. Asterisks (*) indicate significant differences between normoxia and acute hypoxia values; daggers (†) indicate significant differences between AMP- and saline-treated animals ($p < 0.05$).




Figure 3. Naked mole rats do not exhibit a hypoxic change in their air convection requirement (ACR). (a,b) ACR of naked mole rats breathing normoxic or acute hypoxic (7% O₂) gas mixtures and treated with subcutaneous saline (white circles, dashed lines) or AMP (black squares, solid lines) injection. (c) Average RER values from the animals treated in (a,b). Data are mean \pm s.e.m. from $n = 7$ naked mole rats per group. Asterisks (*) indicate significant differences between normoxia and acute hypoxia values; daggers (†) indicate significant differences between AMP- and saline-treated animals ($p < 0.05$).

magnitudes of the HVR and HMR were equal. Similarly, the ACR of AMP-treated naked mole rats was not significantly different from saline-treated naked mole rats when both groups were breathing normoxic air (ACR_{O₂}: 34.1 \pm 10.8;

ACR_{CO₂}: 46.6 \pm 7.8). Conversely, in AMP-treated naked mole rats, the ACR_{O₂} increased by 260% to 88.8 \pm 6.2, whereas the ACR_{CO₂} increased by 167% to 77.3 \pm 9.3 when animals were exposed to 7% O₂ because the HVR was

prevented by AMP treatment, but the HMR was not. These hypoxia-induced changes were reversed upon reoxygenation and indicate that AMP-treated naked mole rats hyperventilated during acute hypoxia (i.e. \dot{V}_E increased relative to metabolic demand because the metabolic rate decreased in hypoxia but \dot{V}_E was unchanged when adenosine receptors were simultaneously blocked).

The RER was not significantly altered in animals breathing 7% O₂ relative to normoxia, or by AMP treatment in either group (figure 3c). Respiratory exchange ratio values were slightly reduced upon recovery, indicating that the animals had not reached steady state, although their ventilatory and metabolic values were not statistically different from pre-hypoxia controls at this time point.

4. Discussion

Naked mole rats are the most hypoxia-tolerant mammals identified. In the present paper, we explore the metabolic, ventilatory and thermal responses of naked mole rats exposed to acute hypoxia, and the potential role for adenosine receptors in mediating these responses. Our study yielded three novel findings. First, naked mole rats undergo robust and reversible metabolic rate and ventilatory depression in acute hypoxia while maintaining T_b . Second; the 70% decrease in metabolic rate during acute hypoxia is matched equally by a 70% decrease in \dot{V}_E . Therefore, \dot{V}_E relative to metabolism (i.e. the ACR) is unchanged during acute hypoxia in naked mole rats, and thus, unique among mammalian responses to hypoxia; naked mole rats do not express a hypoxia-mediated increase in their ACR. Third, systemic antagonism of adenosine receptors prevents the HVR but does not affect the HMR.

(a) Naked mole rats have a robust hypoxic metabolic response

Naked mole rats undergo a profound and reversible HMR when exposed to 7% O₂ for 1 h. To our knowledge, only one other study has examined the effect of hypoxia on metabolic rate in naked mole rats. Nathaniel *et al.* [15] measured metabolism in awake, unrestrained naked mole rats exposed to 3% O₂ for 1 h [15]. These authors reported a significant but very small (8%) hypoxic decrease in metabolic rate. Importantly, the resting metabolic rate reported in their study is approximately 1/10th that observed in our study, and both this difference and also the difference between our measurements of the extent of the HMR (8% and 70%, respectively) are likely to be due to problems with experimental design in the earlier study. Specifically, these authors used a chamber with a volume of approximately 3.5 l, a flow rate of 1 l min⁻¹, and solely reported O₂ consumption. Relative to the small body size of naked mole rats (approx. 45 g) and related low tidal volumes, this large chamber and high flow rate probably saturated the measurement of metabolism in normoxia and hypoxia, resulting in a significant underestimation of the resting metabolic rate and the HMR in the earlier study. Comparatively, our study used a chamber with a considerably smaller volume (approx. 450 ml) and a low flow rate of 110 ml min⁻¹, which provided us greater sensitivity in our measurements of O₂ consumption and CO₂ production. In support of this, our normoxic metabolic rate measurements agree well with other measurements at a similar T_a in this species [32,33].

The cellular mechanism underlying the HMR of naked mole rats has not been examined previously; however, as described in the introduction, several recent studies have demonstrated a central role for adenosinergic signalling in regulating metabolic responses in other small rodents [6–11], although the mechanism via which adenosine receptors modulate metabolism have not been elucidated in any species. Therefore, we also asked whether or not adenosinergic signalling regulates systemic responses to hypoxia in this species. To our surprise, we observed no effect of a general adenosine receptor antagonist (i.e. AMP) on the HMR of the naked mole rat. This indicates that the role of adenosine in mediating the HMR is different in naked mole rats compared with other small mammals. The mechanism that induces and regulates the HMR in naked mole rats requires further study.

(b) Naked mole rats do not exhibit a thermal response to hypoxia

Naked mole rat T_b was approximately 32°C in our study and was not altered by saline or AMP injection, nor by exposure to hypoxia. Animals were held at a T_a of approximately 28°C and thus had only a small thermal range to accommodate hypoxia via metabolic rate depression associated with reductions in T_b . Our findings indicate that naked mole rats do not employ thermal strategies to tolerate hypoxia and that changes in T_b are probably not a component of their HMR. Our findings are in agreement with those of Nathaniel *et al.* [15]. These authors used surgically implanted emitters to record T_b throughout their experiment and reported an average T_b of 28.5 ± 0.2°C in normoxia, and no change in T_b in animals breathing 3% O₂. Similar to our study, these animals were held at a T_a of 28°C, and thus had a minimal range to respond to hypoxia using thermal adaptations. We report a slightly higher T_b than the previous study, but in our experiments all animals remained active and alert, and dug and chewed at the chamber walls throughout the experiment, which may have resulted in a slightly increased T_b . Furthermore, we utilized a smaller chamber with a lower flow rate than the earlier study, which would reduce the rate of animal-generated heat loss from the chamber via thermal convection. To our knowledge, naked mole rats are the only mammals studied to date that exhibit such a robust metabolic depression in acute hypoxia without a concomitant decrease in T_b and entry into a torpor-like state [34]. These unique responses may be due to the warm and consistently hypoxic environment in which naked mole rats live. A high burrow T_a would restrict the scope for thermal adaptations to hypoxia; reducing physical activity in hypoxia would not be a sensible strategy as fresh air is typically only available to the colony through digging new tunnels and vents, which requires physical work. Given these environmental constraints, it appears that naked mole rats have evolved mechanisms of metabolic regulation that are independent of changes in T_b and physical activity.

(c) The naked mole rat hypoxic ventilatory response is mediated by adenosine receptors

Naked mole rats exhibit a robust decrease in \dot{V}_E when exposed to hypoxia for 1 h, which is atypical of mammals [4,34]. Instead, a hypoxia-mediated increase in \dot{V}_E is

consistently observed in other small mammals, and even in other fossorial species, which are typically also tolerant to prolonged hypoxia. For example, degus (*Octodon degus*), coruros (*Spalacopus cyanus*), plateau pikas (*Ochotona curzoniae*) and mole rats (*Spalax ehrenbergi*, a close cousin of naked mole rats) all exhibit increases in \dot{V}_E when in acute hypoxia [13,34–36]. The decrease in \dot{V}_E in hypoxia may represent a beneficial adaptation in naked mole rats since increased breathing, and indeed all other physiological mechanisms aimed at increasing O_2 supply, are energetically expensive responses to hypoxia. In support of this, we observed that many naked mole rats struggled in hypoxia when adenosine receptors were blocked, and in one animal this combination of treatments was lethal, which led us to halt some experiments early in other cases where animals appeared to struggle in hypoxia following AMP injection. Given that naked mole rats are capable of tolerating hypoxia as low as 3% O_2 in a laboratory setting, it is conceivable that they may still exhibit the more typical mammalian HVR of increasing \dot{V}_E at lower inspired O_2 levels; however, the level of hypoxia chosen for our study is similar to the most extreme level of environmental hypoxia experienced in nature by these organisms, and it is notable that other hypoxia-tolerant species exhibit a ‘normal’ HVR of increased breathing within the range of hypoxia they experience in their natural environment (e.g. plateau pika [35]). Nonetheless, further experiments examining the HMR and HVR in naked mole rats at deeper levels of hypoxia would be informative.

Unexpectedly, adenosine receptor antagonism with AMP entirely abolished the decrease in \dot{V}_E in naked mole rats breathing 7% O_2 . This inhibitory effect of adenosine on the HVR is consistent with previous studies in the literature, although the magnitude of its effect is unique to this organism. Adenosine is an inhibitory neurotransmitter that is known to accumulate in the blood during hypoxia and modulate the HVR [37]. Specifically, AMP treatment reduces (but does not abolish) the HVR in awake humans [38], cats [39], rats [40] and lambs [41], and in anaesthetized piglets [42], via modulation of the sensitivity of peripheral chemoreceptors. However, it is notable that in studies in which the effect of AMP on metabolism is examined along with its effects on \dot{V}_E , AMP treatment always inhibits the HMR in addition to the HVR [41,43,44], indicating that the HVR and HMR are coupled in other mammals.

(d) The naked mole rat air convection requirement is not changed in hypoxia

In naked mole rats breathing hypoxic gas mixtures, the magnitudes of the HMR and HVR are equal. Therefore, the ACR is unaltered by hypoxia in this species. To our knowledge, this insensitivity of the ACR to hypoxia is unique among animals

studied to date [4,34]. In addition, we report that AMP injection inhibits hypoxia-mediated ventilatory changes but has no effect on the hypoxia-mediated metabolic changes, indicating that the HMR and HVR are uncoupled by adenosine receptor antagonism in this species. As a result, AMP-treated naked mole rats hyperventilate in hypoxia, as indicated by an increase in their ACR.

Importantly, as the HMR is not altered when the HVR is blocked with AMP, it appears that the HMR is not regulated by O_2 limitation (because AMP blocked the hypoxia-mediated decrease in breathing but not the change in metabolic rate, and thus AMP-treated animals hyperventilate, which should increase the amount of available O_2 at the cellular level and enable a higher metabolic rate). As a result, the increase in metabolic demand associated with increased \dot{V}_E following AMP treatment in hypoxia is not matched by an increase in metabolic rate, leading to a mismatch between O_2 supply and demand and deleterious effects to the organism. In support of this, AMP-treated naked mole rats struggled in hypoxia relative to saline-treated animals, and one naked mole rat died due to hypoxia exposure following AMP injection.

5. Conclusion

We report that naked mole rats, the most hypoxia-tolerant mammal presently identified, undergo a profound metabolic rate depression of 70% when breathing 7% O_2 , which is independent of changes in T_b and matched by an identical decrease in \dot{V}_E . As a result, breathing does not change in hypoxia relative to metabolism, and naked mole rats do not exhibit a hypoxia-mediated change in their ACR. This is a unique response to hypoxia among mammals studied to date. Furthermore, unlike in other mammals, the HMR is insensitive to adenosine receptor inhibition with AMP, whereas the HVR is entirely reversed by AMP. Our findings indicate that naked mole rats may have evolved unique adaptive responses to hypoxia that permit them to decrease metabolic rate independently of changes in T_b and without the benefit of increased O_2 delivery via an energetically costly increase in their ACR. The specific mechanism of this HMR remains unknown, but probably involves cell-level adaptations that enable reduced metabolic rate through reductions in ion pumping, alternative fuel substrate usage or enhanced mitochondrial efficiency [2,3,45].

Ethics statement. All protocols were performed with the approval of the University of British Columbia Animal Care Program and in accordance with the relevant guidelines of the Canadian Council on Animal Care.

Data accessibility. Raw data used in this study can be found at doi:10.5061/dryad.q2h13.

Funding statement. This work was supported by an NSERC Discovery grant to W.K.M. and a Parker B. Francis Foundation PDF to M.E.P.

References

- Buck LT, Pamerter ME. 2006 Adaptive responses of vertebrate neurons to anoxia: matching supply to demand. *Respir. Physiol. Neurobiol.* **154**, 226–240. (doi:10.1016/j.resp.2006.03.004)
- Bickler PE, Buck LT. 2007 Hypoxia tolerance in reptiles, amphibians, and fishes: life with variable oxygen availability. *Annu. Rev. Physiol.* **69**, 145–170. (doi:10.1146/annurev.physiol.69.031905.162529)
- Pamerter ME. 2014 Mitochondria: a multimodal hub of hypoxia tolerance. *Can. J. Zool.* **92**, 569–589. (doi:10.1139/cjz-2013-0247)
- Powell FL, Milsom WK, Mitchell GS. 1998 Time domains of the hypoxic ventilatory response. *Respir. Physiol.* **112**, 123–134. (doi:10.1016/S0034-5687(98)00026-7)
- Barnes BM. 1989 Freeze avoidance in a mammal—body temperatures below 0 degrees C in an Arctic

- hibernator. *Science* **244**, 1593–1595. (doi:10.1126/science.2740905)
6. Tupone D, Madden CJ, Morrison SF. 2013 Central activation of the A1 adenosine receptor (A1AR) induces a hypothermic, torpor-like state in the rat. *J. Neurosci.* **33**, 14 512–14 525. (doi:10.1523/JNEUROSCI.1980-13.2013)
 7. Jinka TR, Toien O, Drew KL. 2011 Season primes the brain in an Arctic hibernator to facilitate entrance into torpor mediated by adenosine A(1) receptors. *J. Neurosci.* **31**, 10 752–10 758. (doi:10.1523/jneurosci.1240-11.2011)
 8. Olson JM, Jinka TR, Larson LK, Danielson JJ, Moore JT, Carpluck J, Drew KL. 2013 Circannual rhythm in body temperature, torpor, and sensitivity to A(1) adenosine receptor agonist in Arctic ground squirrels. *J. Biol. Rhythm.* **28**, 201–207. (doi:10.1177/0748730413490667)
 9. Iliiff BW, Swoap SJ. 2012 Central adenosine receptor signaling is necessary for daily torpor in mice. *Am. J. Physiol.* **303**, R477–R484. (doi:10.1152/ajpregu.00081.2012)
 10. Iliiff BW, Swoap SJ. 2010 Treatment with an adenosine receptor antagonist completely blocks fasting-induced torpor in mice. *FASEB J.* **24**, 1032–1040.
 11. Swoap SJ, Iliiff BW, Le S. 2012 Adenosine, AMP, and daily torpor. In *Living in a seasonal world* (eds T Ruf, C Bieber, W Arnold, E Millesi), pp. 337–339. Vienna, Austria: Springer.
 12. Sherman PW, Jarvis JU, Alexander RD. 1991 *The biology of the naked mole-rat*. Princeton, NJ: Princeton University Press.
 13. Arieli R, Ar A. 1979 Ventilation of a fossorial mammal (*Spalax ehrenbergi*) in hypoxic and hypercapnic conditions. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* **47**, 1011–1017.
 14. Shams I, Avivi A, Nevo E. 2005 Oxygen and carbon dioxide fluctuations in burrows of subterranean blind mole rats indicate tolerance to hypoxic-hypercapnic stresses. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **142**, 376–382. (doi:10.1016/j.cbpa.2005.09.003)
 15. Nathaniel TI, Otukonyong E, Abdellatif A, Soyinka JO. 2012 Effect of hypoxia on metabolic rate, core body temperature, and c-fos expression in the naked mole rat. *Int. J. Dev. Neurosci.* **30**, 539–544. (doi:10.1016/j.ijdevneu.2012.04.004)
 16. Johansen K, Lykkeboe G, Weber RE, Maloij GM. 1976 Blood respiratory properties in the naked mole rat *Heterocephalus glaber*, a mammal of low body temperature. *Respir. Physiol.* **28**, 303–314. (doi:10.1016/0034-5687(76)90025-6)
 17. Buffenstein R, Yahav S. 1991 Is the naked mole-rat, *Heterocephalus glaber*, a poikilothermic or poorly thermoregulating endothermic mammal? *J. Therm. Biol.* **16**, 227–232. (doi:10.1016/0306-4565(91)90030-6)
 18. Peterson BL, Larson J, Buffenstein R, Park TJ, Fall CP. 2012 Blunted neuronal calcium response to hypoxia in naked mole-rat hippocampus. *PLoS ONE* **7**, e31568. (doi:10.1371/journal.pone.0031568)
 19. Peterson BL, Park TJ, Larson J. 2012 Adult naked mole-rat brain retains the NMDA receptor subunit GluN2D associated with hypoxia tolerance in neonatal mammals. *Neurosci. Lett.* **506**, 342–345. (doi:10.1016/j.neulet.2011.11.042)
 20. Nathaniel TI, Saras A, Umehiri FE, Olajuyigbe F. 2009 Tolerance to oxygen nutrient deprivation in the hippocampal slices of the naked mole rats. *J. Integr. Neurosci.* **8**, 123–136. (doi:10.1142/S0219635209002149)
 21. Bickler PE, Buck LT. 1998 Adaptations of vertebrate neurons to hypoxia and anoxia: maintaining critical Ca²⁺ concentrations. *J. Exp. Biol.* **201**, 1141–1152.
 22. Aranda JV, Turmen T. 1979 Methylxanthines in apnea of prematurity. *Clin. Perinatol.* **6**, 87–108.
 23. Tanikawa K, Matsumoto Y, Matsuzaki T, Matsumoto M, Fukuoka M, Noguchi S, Goshima T. 1999 Population pharmacokinetic analysis of theophylline: relationship between serum concentrations and clinical effects in therapeutic drug monitoring. *Yakugaku Zasshi* **119**, 861–867.
 24. Bach JE, Kukanich B, Papich MG, McKiernan BC. 2004 Evaluation of the bioavailability and pharmacokinetics of two extended-release theophylline formulations in dogs. *J. Am. Vet. Med. Assoc.* **224**, 1113–1119. (doi:10.2460/javma.2004.224.1113)
 25. Yin B, Li J, Wang SL, Yan SY, Sui C, Wan SJ, Liu F, Wan RZ. In press. Studies on pharmacokinetics and bioavailability of aminophylline in partridge chickens. *Eur. J. Drug Metab. Pharmacokinet.* (doi:10.1007/s13318-014-0225-6)
 26. Asheim P, Spigset O, Aasarod K, Walstad RA, Uggen PE, Zahlsen K, Aadahl P. 2008 Pharmacokinetics of intraperitoneally instilled aminophylline, terbutaline and tobramycin in pigs. *Acta Anaesthesiol. Scand.* **52**, 243–248. (doi:10.1111/j.1399-6576.2007.01535.x)
 27. Jacky JP. 1978 A plethysmograph for long-term measurements of ventilation in unrestrained animals. *J. Appl. Physiol.* **45**, 644–647.
 28. Drorbaugh JE, Fenn WO. 1955 A barometric method for measuring ventilation in newborn infants. *Pediatrics* **16**, 81–87.
 29. Mearthur MD, Milsom WK. 1991 Ventilation and respiratory sensitivity of euthermic Columbian and golden-mantled ground-squirrels (*Spermophilus columbianus* and *Spermophilus lateralis*) during the Summer and Winter. *Physiol. Zool.* **64**, 921–939.
 30. Frayn KN. 1983 Calculation of substrate oxidation rates *in vivo* from gaseous exchange. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* **55**, 628–634.
 31. Koupenova M, Ravid K. In press. Adenosine, adenosine receptors and their role in glucose homeostasis and lipid metabolism. *J. Cell Physiol.* (doi:10.1002/jcp.24352)
 32. Goldman BD, Goldman SL, Lanz T, Magaurin A, Maurice A. 1999 Factors influencing metabolic rate in naked mole-rats (*Heterocephalus glaber*). *Physiol. Behav.* **66**, 447–459. (doi:10.1016/S0031-9384(98) 00306-0)
 33. O'Connor TP, Lee A, Jarvis JU, Buffenstein R. 2002 Prolonged longevity in naked mole-rats: age-related changes in metabolism, body composition and gastrointestinal function. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **133**, 835–842. (doi:10.1016/S1095-6433(02)00198-8)
 34. Frappell P, Lanthier C, Baudinette RV, Mortola JP. 1992 Metabolism and ventilation in acute hypoxia—a comparative analysis in small mammalian species. *Am. J. Physiol.* **262**, R1040–R1046.
 35. Pichon A, Zhenzhong B, Favret F, Jin G, Shufeng H, Marchant D, Richalet JP, Ge RL. 2009 Long-term ventilatory adaptation and ventilatory response to hypoxia in plateau pika (*Ochotona curzoniae*): role of nNOS and dopamine. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **297**, R978–987. (doi:10.1152/ajpregu.00108.2009)
 36. Tomasco IH, Del Rio R, Iturriaga R, Bozinovic F. 2010 Comparative respiratory strategies of subterranean and fossorial octodontid rodents to cope with hypoxic and hypercapnic atmospheres. *J. Comp. Physiol. B.* **180**, 877–884. (doi:10.1007/s00360-010-0465-y)
 37. Drumm D, Hoefler M, Juhasz J, Huszar E, Sybrecht GW. 2004 Plasma adenosine during investigation of hypoxic ventilatory response. *Sleep Breath* **8**, 31–41. (doi:10.1007/s11325-004-0031-5)
 38. Easton PA, Anthonisen NR. 1988 Ventilatory response to sustained hypoxia after pretreatment with aminophylline. *J. Appl. Physiol.* **64**, 1445–1450.
 39. Long WQ, Anthonisen NR. 1994 Aminophylline partially blocks ventilatory depression with hypoxia in the awake cat. *Can. J. Physiol. Pharmacol.* **72**, 673–678. (doi:10.1139/y94-095)
 40. Maskrey M, Westwood KJ. 1994 Dual effect of aminophylline on the ventilatory response to hypoxia in the rat. *J. Basic Clin. Physiol. Pharmacol.* **5**, 227–238. (doi:10.1515/JBCPP.1994.5.3-4.227)
 41. Koos BJ, Kawasaki Y, Kim YH, Bohorquez F. 2005 Adenosine A2A-receptor blockade abolishes the roll-off respiratory response to hypoxia in awake lambs. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **288**, R1185–1194. (doi:10.1152/ajpregu.00723.2004)
 42. Darnall Jr RA. 1985 Aminophylline reduces hypoxic ventilatory depression: possible role of adenosine. *Pediatr. Res.* **19**, 706–710. (doi:10.1203/00006450-198507000-00014)
 43. Barros RC, Branco LG, Carnio EC. 2006 Respiratory and body temperature modulation by adenosine A1 receptors in the anteroventral preoptic region during normoxia and hypoxia. *Respir. Physiol. Neurobiol.* **153**, 115–125. (doi:10.1016/j.resp.2005.09.013)
 44. Maxwell DL, Fuller RW, Noland KB, Dixon CM, Hughes JM. 1986 Effects of adenosine on ventilatory responses to hypoxia and hypercapnia in humans. *J. Appl. Physiol.* **61**, 1762–1766.
 45. Cheviron ZA, Bachman GC, Connaty AD, McClelland GB, Storz JF. 2012 Regulatory changes contribute to the adaptive enhancement of thermogenic capacity in high-altitude deer mice. *Proc. Natl Acad. Sci. USA* **109**, 8635–8640. (doi:10.1073/pnas.1120523109)