



## Research

**Cite this article:** Sun B-J, Li T, Mu Y, McGlashan JK, Georges A, Shine R, Du W-G. 2016 Thyroid hormone modulates offspring sex ratio in a turtle with temperature-dependent sex determination. *Proc. R. Soc. B* **283**: 20161206.  
<http://dx.doi.org/10.1098/rspb.2016.1206>

Received: 1 June 2016

Accepted: 29 September 2016

**Subject Areas:**

ecology, physiology

**Keywords:**

embryonic development, sex determination, thyroid hormone, turtle

**Author for correspondence:**

Wei-Guo Du

e-mail: [duweigu@ioz.ac.cn](mailto:duweigu@ioz.ac.cn)

<sup>†</sup>These authors contributed equally to this study.

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.3518949>.

# Thyroid hormone modulates offspring sex ratio in a turtle with temperature-dependent sex determination

Bao-Jun Sun<sup>1,†</sup>, Teng Li<sup>1,†</sup>, Yi Mu<sup>1,†</sup>, Jessica K. McGlashan<sup>2</sup>, Arthur Georges<sup>3</sup>, Richard Shine<sup>4</sup> and Wei-Guo Du<sup>1</sup>

<sup>1</sup>Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing, 100101, People's Republic of China

<sup>2</sup>Science and Health Hawkesbury Institute, University of Western Sydney, New South Wales 2751, Australia

<sup>3</sup>Institute for Applied Ecology, University of Canberra, Australian Capital Territory 2601, Australia

<sup>4</sup>School of Life and Environmental Sciences A08, University of Sydney, New South Wales 2006, Australia

B-JS, 0000-0002-7318-6059; RS, 0000-0001-7529-5657; W-GD, 0000-0002-1868-5664

The adaptive significance of temperature-dependent sex determination (TSD) has attracted a great deal of research, but the underlying mechanisms by which temperature determines the sex of a developing embryo remain poorly understood. Here, we manipulated the level of a thyroid hormone (TH), triiodothyronine (T<sub>3</sub>), during embryonic development (by adding excess T<sub>3</sub> to the eggs of the red-eared slider turtle *Trachemys scripta*, a reptile with TSD), to test two competing hypotheses on the proximate basis for TSD: the *developmental rate hypothesis* versus the *hormone hypothesis*. Exogenous TH accelerated embryonic heart rate (and hence metabolic rate), developmental rate, and rates of early post-hatching growth. More importantly, hyperthyroid conditions depressed expression of *Cyp19a1* (the gene encoding for aromatase) and levels of oestradiol, and induced more male offspring. This result is contrary to the direction of sex-ratio shift predicted by the *developmental rate hypothesis*, but consistent with that predicted by the *hormone hypothesis*. Our results suggest an important role for THs in regulating sex steroid hormones, and therefore, in affecting gonadal sex differentiation in TSD reptiles. Our study has implications for the conservation of TSD reptiles in the context of global change because environmental contaminants may disrupt the activity of THs, and thereby affect offspring sex in TSD reptiles.

## 1. Introduction

Despite strong genetic similarity, males and females within a population often differ dramatically in a diverse suite of phenotypic traits encompassing behaviour, physiology, and morphology [1–3]. Thus, the gene regulatory pathways that determine whether an individual develops as a male or a female are fundamental to a broad range of processes in ecology and evolution [4–6]. Sex is genotypically determined in mammals, birds, snakes, most amphibians, and many other vertebrates; that is, an offspring's sex depends upon the complement of sex chromosomes that it receives from its parents at the time of conception. By contrast, sex in many ectothermic vertebrates is determined by a remarkable range of other systems that include environmental triggers. For example, nest temperature determines offspring sex in all crocodylians [7], many turtles, and lizards [8,9], and the sole extant rhynchocephalian [10]. Both the ultimate (adaptive) and proximate foundations of temperature-dependent sex determination (TSD) have attracted a great deal of research [11–13], but the underlying mechanisms by which temperature determines the sex of a developing embryo remain poorly understood, and a major challenge for evolutionary biology [14–16].

In this paper, we focus on two competing hypotheses regarding the proximate basis for TSD. The first, which we call the *developmental rate hypothesis*, posits that sex is determined by embryonic development rate rather than

incubation temperature *per se* [17]. In support of this idea, an early study on crocodiles found that offspring sex is more closely related to embryonic development rate during the first half of incubation than it is to incubation temperature [18]. Experimental tests of this hypothesis have been less encouraging. Hatchling sex in turtles is unaffected by manipulations that induce shifts in developmental rate (changes in hydric or oxygen incubation environments [19,20]), nor is sex associated with intraspecific variations in genome size that influence rates of cell division and growth [21].

The second and more broadly accepted hypothesis, which we refer to as the *hormone hypothesis*, posits a central role for levels of sex steroid hormones (androgens and oestrogens), moderated by as-yet-unknown temperature-sensitive genes that direct development into either a male or female trajectory [22]. Support for a central role of sex hormones is provided by experiments that block aromatase (which converts testosterone to oestradiol ( $E_2$ )) to influence the outcome of the female/male determining cascade [23,24]; and by the ability of exogenous  $E_2$  to override the effect of temperature on gonadal sex. Applying aromatase inhibitors can generate male offspring at female-producing temperatures, and applying exogenous  $E_2$  can generate female offspring at male-producing temperatures [13,25,26]. Nonetheless, the mechanism by which temperature influences levels of sex steroid hormones, and thus offspring sex, remains unclear.

Other hormones play a critical role both in upregulating or downregulating the level of expression of genes affecting androgens and oestrogens, and in influencing the overall rate of development of the embryo. One such class of hormones, thyroid hormones (THs), regulate metabolic rate during the early growth and development of vertebrates [27–29]. THs control rates of organ differentiation (and thus the pace of embryonic and larval development) in fish, amphibians, reptiles, and birds [30–33]. TH also influences the establishment of gonadal sex in fish, amphibians, and mammals. For example, treatment with goitrogenic compounds or perchlorate (that inhibit TH synthesis) increases the proportion of females among zebrafish (*Danio rerio*) and anuran embryos, whereas application of TH yields male-skewed sex ratios [34–36]. Uniquely, then, TH not only stimulates developmental rate, but also affects gonadal sex. Experiments that modify TH levels thus can directly test contrasting predictions from the *developmental rate hypothesis* and the *hormone hypothesis*.

In this study, we manipulated the level of TH (triiodothyronine,  $T_3$ ) during embryonic development by adding excess  $T_3$  to the eggs of the red-eared slider turtle (*Trachemys scripta*), a reptile with TSD. The eggs were incubated at three constant temperatures (28°C, 29°C, and 30°C) around the pivotal temperature of 29°C [37] to explore the influence of  $T_3$  on developmental rate, levels of expression of *Cyp19a1* (Cytochrome P450 Family 19 Subfamily A Member 1, encoding for aromatase), oestradiol levels in embryos and offspring sex. Based on the *developmental rate hypothesis*, we expect  $T_3$  treated eggs to produce more females, because faster developmental rates (as generated by higher temperatures, and also by  $T_3$  administration) result in females in this species [37]. Alternatively, if  $T_3$  directly affects oestrogen levels, consistent with the *hormone hypothesis*, we predict lower levels of expression of aromatase and, as a result, lower levels of oestradiol and a male-biased sex ratio in the  $T_3$  treatment (because elevated levels of  $T_3$  and its precursor thyroxine  $T_4$  induce more male offspring [35,36]).

## 2. Material and methods

### (a) Study species

The red-eared slider turtle (*T. scripta*) is widespread globally, due to a flourishing pet trade and generalist ecological niche [38]. *Trachemys scripta* exhibits Type I TSD, whereby 100% females are produced at a constant incubation temperature greater than 30°C and 100% males at less than 26°C, with a pivotal temperature of 29°C [37,39]. Females lay on average 12 eggs that weigh between 9 and 11 g and neonates have been recorded to overwinter in the nest and emerge the following spring (in cool-climate areas: [38,40]).

### (b) Egg collection and incubation

In April 2015, we collected freshly laid eggs from a private hatchery in Haikou, Hainan province. Seventy-five clutches of fertilized eggs ( $N = 577$ ) were weighed to the nearest 1 mg on a Mettler balance, and each was individually incubated in an 80 ml jar half-filled with moist vermiculite (water tension  $-220$  kPa) [41]. Each jar was assigned to either a treatment group ( $T_3$  application) or control group (ethanol (EtOH)). Both treatment and control jars were distributed among three incubators (KB240, Binder, Germany) set at 28°C, 29°C, and 30°C. Those temperatures were predicted to result in male-biased sex ratios, even sex ratios, and female-biased sex ratios, respectively [39]. Jars were weighed and rehydrated weekly to account for evaporation and maintain water potential of vermiculite. To counteract potential thermal gradients in the incubator, jars were rotated weekly within chambers. Temperatures within each incubator were monitored using Thermocron iButton temperature loggers (two DS1921 iButtons in separate jars, MAXIM Integrated Products Ltd., USA), so that we could record the exact thermal environments experienced by the eggs.

### (c) Application of thyroid hormone

Eggs in the treatment group were exogenously treated with the TH 3,3',5-triiodo-L-thyronine ( $T_3$ ; Sigma-Aldrich, USA) dissolved in absolute ethanol (EtOH, Sigma-Aldrich, USA)—4.0  $\mu\text{g}$   $T_3$  was dissolved in 10.0  $\mu\text{l}$  of EtOH to yield a concentration of 0.4  $\mu\text{g}$   $\mu\text{l}^{-1}$  [42]. Eggs in the control group were treated with EtOH only. The dosage was based on the dosages applied to *Chelydra serpentina* and *Emydera macquarii* embryos at embryonic stages 18–19 [42,43], after estimating and correcting for estimated embryo size (crown to rump length) of *T. scripta* embryo size at the time of application [44,45]. We then conducted a preliminary experiment on  $T_3$  dosage effects to determine the optimal dosage, which was 3  $\mu\text{l}$  of  $T_3$  solution (0.4  $\mu\text{g}$   $\mu\text{l}^{-1}$ ) (electronic supplementary material, figure S1).

Every 3 days through the incubation period, we collected three eggs from each incubator and dissected them to determine the stage of embryonic development. When embryos reached stage 15 according to the classification scheme of Greenbaum [45], the beginning of the thermosensitive period of TSD [46], each egg was given a single dose of 3  $\mu\text{l}$  of either the treatment (0.4  $\mu\text{g}$   $T_3$   $\mu\text{l}^{-1}$  EtOH) or control (EtOH) solutions, applied to the top surface of each egg using a pipette.

### (d) Aromatase and oestradiol during the thermosensitive period of temperature-dependent sex determination

After the application of  $T_3$ , we dissected three eggs every other day from each treatment to determine the developmental stage of embryos. Embryos at stages 17, 19, and 21 were collected to determine the expression of aromatase and measure levels of oestradiol.

For aromatase, we collected a total of 90 embryos (five embryos at each stage from each treatment). The Adrenal Kidney Gonad Complex (AKG) of each embryo was separated immediately on dissection and frozen in liquid nitrogen. RNA was extracted with TRIzol reagent (Invitrogen, Carlsbad, CA, USA) following the manufacturer's protocols. The quality and quantity of each total RNA sample was assessed through spectrophotometer reading at 260 nm and 280 nm. The RNA sample was then treated with DNase I to eliminate genomic DNA contamination and single-stranded complementary DNA (cDNA) was reverse-transcribed using an RT reagent kit with genomic DNA (gDNA) eraser (Takara Bio Inc., CA, USA). After reverse transcription, we mixed the cDNA of five embryos at each stage from each treatment together to obtain enough cDNA for polymerase chain reaction (PCR). Relative levels of *Cyp19a1* expression were quantified using SYBR Green (Hercules, CA, USA) following the manufacturer's protocol and Roche LightCycler 480 real-time PCR cyler. Each reaction was performed in triplicate and normalized to constitutive expression of peptidylprolyl isomerase (PPI). Primers used to assay gene expression were designed according to Ramsey *et al.* [47]. The qPCR cycling conditions were 95°C for 5 min, 40 cycles of 95°C for 30 s, 59°C for 20 s, 72°C for 20 s, and then melt curve from 55 to 95°C. Relative gene expression was calculated from the equation  $2^{-\Delta\Delta CT}$ .

For measuring levels of oestradiol, we collected 125 embryos (six to eight embryos at each stage from each treatment). Each embryo was homogenized with a glass homogenizer in ice cold phosphate buffer saline (PBS). The suspensions were then centrifuged at 13 000 r.p.m. for 10 min. The volume of each supernatant was only large enough for us to assay oestradiol levels with radioimmunoassay. Even so, 11 embryos were too small for us to detect the oestradiol level accurately, and data on these individuals were excluded from subsequent analyses.

### (e) Embryonic heart rate

We measured heart rates of 15 eggs from each treatment 24 h after  $T_3$  application, in order to infer the metabolic rate of embryos [48]. For this purpose, eggs were individually removed from the incubation container and immediately placed in a Buddy digital egg monitor system (Avian Biotech, UK) in complete darkness for a maximum of 2 min [48,49]. Eggs were not disturbed during the recording period. If any embryonic movement was detected, we waited for movement to cease and heart rate to stabilize before recording [49].

### (f) Incubation duration and hatching success

Towards the end of incubation, eggs were inspected twice daily for signs of pipping (initial break of egg with caruncle) [50]. The duration of incubation was measured for each egg as the number of days from oviposition to pipping. Hatching success was recorded as the number of neonates that hatched as a percentage of all eggs assigned to the respective treatment.

### (g) Hatchling traits

Directly after the eggs hatched, we measured maximum carapace length, width, and height, and mass of each hatchling. The young turtles were each individually housed in plastic containers (150 × 100 × 60 mm) for the next three months. The containers were set in a temperature-controlled room at  $28 \pm 0.5^\circ\text{C}$ , with a 12 light (L):12 dark (D) photoperiod. Sufficient commercial food was provided for hatchlings every day. Hatchling size and growth rate were recorded twice (at one and three months of age) to measure the effects of TH and incubation temperature on post-hatching development. Hatchlings were sexed at three months of age, when the sex of the turtles can be reliably identified by their secondary sexual characteristics [51].

### (h) Data analysis

We used the Kolmogorov–Smirnov test and Levene's test to evaluate the normality of distributions and homogeneity of variances for all data.  $\chi^2$ -tests were used to explore the effects of incubation temperature and  $T_3$  on hatching success and hatchling sex ratio. Factorial ANOVA was used to detect the effect of temperature and  $T_3$  on embryonic heart rate and incubation duration, as well as the effect of temperature,  $T_3$ , and developmental stage on the expression of aromatase and levels of oestradiol. Two-way ANCOVA was used to analyse the effects of temperature and  $T_3$  on hatchling sizes (with initial egg mass as the covariate) and growth rates (with hatchling carapace length as the covariate). Post hoc Tukey–Kramer honest significant difference (HSD) tests were carried out to determine the locations of among-treatment differences.

## 3. Results

Expression of aromatase in embryos increased with incubation temperature ( $F_{2,36} = 72.1$ ,  $p < 0.00001$ ) and as the embryos developed ( $F_{2,36} = 135.7$ ,  $p < 0.00001$ ), but was depressed by  $T_3$  application ( $F_{1,36} = 23.0$ ,  $p < 0.0001$ ; figure 1). Oestradiol levels in embryos increased as the embryos developed ( $F_{2,96} = 36.2$ ,  $p < 0.00001$ ), were not affected by incubation temperature ( $F_{2,96} = 2.9$ ,  $p = 0.061$ ), but were depressed by  $T_3$  application ( $F_{1,96} = 27.1$ ,  $p < 0.0001$ ; figure 2).


Incubation temperature accelerated embryogenesis: it increased embryonic heart rates and decreased incubation durations (table 1; figure 3). Exogenous TH ( $T_3$ ) also increased embryonic heart rates (by 4–8%: figure 3a), and shortened incubation duration of turtle eggs (by 4–7%: figure 3b, table 1). The effect of  $T_3$  on incubation duration depended on incubation temperature (i.e. significant interaction between  $T_3$  treatment and incubation temperature: table 1). The effect of  $T_3$  was greater at lower incubation temperatures (figure 3).

Hatching success of turtle eggs was high (approx. 90%), and was not significantly affected by either  $T_3$  (28°C:  $\chi^2 = 0.33$ , d.f. = 1,  $p = 0.57$ ; 29°C:  $\chi^2 = 0.78$ , d.f. = 1,  $p = 0.38$ ; 30°C:  $\chi^2 = 0.12$ , d.f. = 1,  $p = 0.73$ ) or incubation temperature ( $T_3$  treatment:  $\chi^2 = 1.388$ , d.f. = 2,  $p = 0.50$ ; control:  $\chi^2 = 0.084$ , d.f. = 2,  $p = 0.96$ ; figure 4).

The sex ratio of hatchlings was affected by exogenous  $T_3$  when eggs were incubated at 28°C ( $\chi^2 = 6.35$ , d.f. = 1,  $p = 0.01$ ) and 29°C ( $\chi^2 = 6.76$ , d.f. = 1,  $p < 0.01$ ), but not at 30°C ( $\chi^2 = 2.65$ , d.f. = 1,  $p = 0.10$ ). Generally,  $T_3$  caused a male-biased sex ratio relative to the control group (figure 5). Hatchling sex was also affected by incubation temperature ( $T_3$  treatment:  $\chi^2 = 94.8$ , d.f. = 2,  $p < 0.001$ ; control:  $\chi^2 = 65.64$ , d.f. = 2,  $p < 0.001$ ). Hatchling sex ratios were male-biased at 28°C and 29°C, but female-biased at 30°C (figure 5).

$T_3$  treatment, incubation temperatures, and the interaction between these two factors also affected hatchling size (carapace length, width, and height) and mass (table 1). Hatchlings from the  $T_3$  treatment were smaller than their counterparts from the control group. In the  $T_3$  treatment, hatchlings from 28°C were smaller than those from 29°C and 30°C, whereas hatchling size was not significantly affected by incubation temperature in the control group (electronic supplementary material, figure S2).

During the first month of life, a hatchling's growth was affected by both the  $T_3$  treatment and incubation temperature under which it had developed as an embryo (table 1). Hatchlings from the  $T_3$  treatment grew more rapidly than did those from the control groups and hatchlings from 28°C grew more




**Figure 1.** Effects of triiodothyronine ( $T_3$ ) treatment on expression of aromatase in different developmental stages of *Trachemys scripta* embryos incubated at (a) 28°C, (b) 29°C, and (c) 30°C. Data are expressed as mean  $\pm$  standard error, and the sample sizes are all three. The asterisk indicates a significant difference between control and  $T_3$  application. (Online version in colour.)

rapidly than did those from 29°C and 30°C (electronic supplementary material, figure S3). By contrast, rates of growth by hatchlings during the second and third months of life were affected by incubation temperature but not by  $T_3$  treatment (table 1). Hatchlings from 28°C grew more rapidly than did those from 29°C and 30°C (electronic supplementary material, figure S4).

#### 4. Discussion

Exogenous TH ( $T_3$ ) accelerated embryonic heart rate (and hence metabolic rate), developmental rate, and rates of early



**Figure 2.** Effects of triiodothyronine ( $T_3$ ) treatment on oestradiol levels in different developmental stages of *Trachemys scripta* embryos incubated at (a) 28°C, (b) 29°C, and (c) 30°C. Data are expressed as mean  $\pm$  standard error, and sample sizes are shown above the error bars. The asterisk indicates a significant difference between control and  $T_3$  application. (Online version in colour.)

post-hatching growth. More importantly, the administration of  $T_3$  during embryonic development depressed expression of aromatase and levels of oestradiol, and induced more male offspring in *T. scripta*. This result is contrary to the direction of sex-ratio shift predicted by the *developmental rate hypothesis*, but consistent with that predicted by the *hormone hypothesis*.

THs play critical roles in metabolism, differentiation, development, and growth [31,32,52–54]. Hyperthyroid conditions enhance oxygen consumption and metabolic rate, and thereby affect growth and development [54]. Consistent with such stimulatory effects, our study indicated that application of exogenous  $T_3$  increased the heart rate (and hence metabolic rate) of embryonic turtles. In addition, exogenous

**Table 1.** Two-way ANOVA or ANCOVA results for embryonic heart rate, incubation length, hatchling sizes, and hatchling growth in *Trachemys scripta*. Growth 1, growth during the first month; growth 2, growth during the subsequent two months. The significant differences are shown in bold.


variable	T <sub>3</sub> treatment	incubation temperature	interaction
heart rate	$F_{1,84} = 55.9$ , $p < 0.00001$	$F_{2,84} = 79.9$ , $p < 0.00001$	$F_{2,84} = 2.2$ , $p = 0.12$
incubation length	$F_{1,272} = 147.8$ , $p < 0.00001$	$F_{2,272} = 361.7$ , $p < 0.00001$	$F_{2,272} = 6.6$ , $p < 0.01$
carapace length (CL)	$F_{1,271} = 23.9$ , $p < 0.00001$	$F_{2,271} = 3.9$ , $p = 0.02$	$F_{2,271} = 1.5$ , $p = 0.24$
carapace width (CW)	$F_{1,271} = 14.4$ , $p < 0.001$	$F_{2,271} = 15.2$ , $p < 0.00001$	$F_{2,271} = 2.6$ , $p = 0.07$
carapace height (CH)	$F_{1,271} = 17.2$ , $p < 0.0001$	$F_{2,271} = 35.8$ , $p < 0.00001$	$F_{2,271} = 11.5$ , $p < 0.001$
body mass (BM)	$F_{1,271} = 26.3$ , $p < 0.0001$	$F_{2,271} = 4.0$ , $p = 0.02$	$F_{2,271} = 7.6$ , $p < 0.001$
growth 1 in CL	$F_{1,268} = 16.2$ , $p < 0.0001$	$F_{2,268} = 1.5$ , $p = 0.22$	$F_{2,268} = 3.5$ , $p = 0.03$
growth 1 in CW	$F_{1,268} = 6.6$ , $p = 0.01$	$F_{2,268} = 5.0$ , $p < 0.01$	$F_{2,268} = 10.8$ , $p < 0.0001$
growth 1 in CH	$F_{1,268} = 0.34$ , $p = 0.56$	$F_{2,268} = 48.3$ , $p < 0.0001$	$F_{2,268} = 10.8$ , $p < 0.0001$
growth 1 in BM	$F_{1,268} = 9.1$ , $p < 0.01$	$F_{2,268} = 28.3$ , $p < 0.0001$	$F_{2,268} = 0.38$ , $p = 0.68$
growth 2 in CL	$F_{1,268} = 1.2$ , $p = 0.27$	$F_{2,268} = 7.1$ , $p = 0.001$	$F_{2,268} = 2.1$ , $p = 0.12$
growth 2 in CW	$F_{1,268} = 0.9$ , $p = 0.34$	$F_{2,268} = 4.9$ , $p < 0.01$	$F_{2,268} = 1.7$ , $p = 0.19$
growth 2 in CH	$F_{1,268} = 3.4$ , $p = 0.07$	$F_{2,268} = 12.8$ , $p < 0.0001$	$F_{2,268} = 3.3$ , $p = 0.04$
growth 2 in BM	$F_{1,268} = 1.4$ , $p = 0.24$	$F_{2,268} = 8.9$ , $p < 0.001$	$F_{2,268} = 2.2$ , $p = 0.11$

T<sub>3</sub> reduced the incubation period of turtle eggs (figure 3). The reduced incubation period may be owing to faster developmental rate, or because the final hatchling size is smaller (and thus, can be achieved more rapidly). In *E. macquarii*, exogenous application of T<sub>3</sub> reduced the incubation period of neonates with no cost to development, suggesting that embryonic development was complete [43]. In zebrafish, an experimental reduction in levels of endogenous TH decreased body lengths of developing embryos, an effect that was not reversed by exogenous application of T<sub>4</sub>, the precursor to T<sub>3</sub> [55]. If the same phenomenon occurs in turtle embryos, our application of exogenous T<sub>3</sub> may have induced a corresponding reduction in the embryo's production of its own T<sub>4</sub> thyroxin, in turn leading to reduced hatchling size. Young turtles from the T<sub>3</sub>-treated eggs grew faster than the controls, indicating a carry-over effect of embryonic T<sub>3</sub> treatment. However, this effect was relatively brief (only seen in the first month after hatching).


Although the close relationship between hatchling sex ratio and developmental rate in crocodiles supports the developmental rate hypothesis [17,18], manipulative studies on turtles have failed to support the idea [19,20,56]. First, a change in developmental rate (induced by incubation

conditions independent of temperature) did not modify the sex ratio of hatchlings [19,20]. Second, Georges *et al.* [56] manipulated the sex ratio of a turtle from 100% male to 100% female by modifying diel schedules of thermal variation while holding overall development rate (incubation duration) constant. Third, changes in developmental rate due to hormones (this study) or thermal conditions [57,58] have been associated with biased sex ratios, but in the opposite direction to that predicted by the developmental rate hypothesis.

Our study provides the first evidence that TH affects the offspring sex ratio in TSD reptiles, as it is known to do in some teleosts and amphibians [34–36]. Generally, hyperthyroid conditions during early development produce more male offspring in vertebrates, whereas hypothyroid conditions produce more female offspring [36,55]. For example, perchlorate, a TH synthesis inhibitor, skewed the sex ratio of zebrafish towards female, but yielded male-skewed sex ratios when co-treated with exogenous TH [36]. Our results suggest that TH affects sex-determining pathways in TSD reptiles (figures 1,2,5). However, we caution that our work is based only on a single species, and studies in fish have




**Figure 3.** Effects of triiodothyronine ( $T_3$ ) treatment on embryonic heart rate (a) and incubation duration (b) in *Trachemys scripta* eggs incubated at different temperatures. Data are expressed as mean  $\pm$  standard error. Numbers above the error bar are sample sizes. (Online version in colour.)



**Figure 4.** Hatching success of *Trachemys scripta* eggs treated with triiodothyronine ( $T_3$ ) and incubated at different temperatures. Numbers in the bars are sample sizes. (Online version in colour.)

reported contradictory results. For example, perchlorate did not affect the gonadal sex phenotype of zebrafish [59], or induce male-skewed sex ratios in three-spined sticklebacks (*Gasterosteus aculeatus*) [60]. The broader question of how TH affects sex-determining mechanisms merits further studies in a wide range of vertebrate species.

The underlying mechanisms by which TH affects sex determination in TSD reptiles may be associated with interactions between TH and the sex steroid hormones (androgen and oestrogen) that appear to play a critical role in determining



**Figure 5.** Effects of triiodothyronine ( $T_3$ ) treatment on the sex ratio of *Trachemys scripta* hatchlings from different incubation temperatures. Numbers in the bars are number of males produced out of sample size. (Online version in colour.)

gonadal sex of TSD reptiles [55,61]. TH might affect sex-determining pathways in several ways. First, TH may affect the activity of gonadal aromatase [27,36]. Aromatase is a key factor in the sex determination of vertebrates; experimental inhibition of aromatase activity reduces oestrogen production and, therefore, generates male-biased sex ratios in species with both genotypic and TSD [22,62,63]. Likewise, hyperthyroid conditions may reduce aromatase activity and thus generate male-biased gonadal sex ratios [36]. Our results on the expression of aromatase support the hypothesis that hyperthyroid conditions can depress aromatase activity (figure 1), which may block the conversion of testosterone to oestradiol and, therefore, promote the male regulatory cascade at the expense of the female cascade [23,24]. Second, TH may upregulate androgen receptors and downregulate oestrogen receptors [61,64,65], again promoting the male regulatory cascade. Unfortunately, we were not able to test this hypothesis in this study. Future studies on how TH affects the regulation of these receptors would deepen our understanding of the physiological mechanism by which hyperthyroid conditions produce more male offspring in reptiles.

Sex steroid hormones play critical roles in determining gonadal sex in TSD reptiles, and thus have been the focus of most studies on physiological pathways underlying TSD in these animals [66]. However, we know little about interactions between sex steroid hormones and other important hormones in determining the sexual phenotype of TSD reptiles. Our results suggest an important role for TH in regulating sex steroid hormones, and therefore, in affecting gonadal sex differentiation in TSD reptiles. For example, application of exogenous  $T_3$  inhibited the activity of aromatase and consequently decreased the level of oestradiol in *T. scripta* (figure 2), thereby inducing male offspring at female-producing temperatures (see also [13,25]). Our study not only highlights the importance of studying interactions among different hormones when we attempt to elucidate physiological mechanisms underlying the TSD in reptiles, but also has implications for the conservation of TSD reptiles under the context of global change. For example, environmental contaminants like perchlorate may disrupt the activity of TH, and thereby affect the sex of TSD reptiles.

**Ethics.** This research was performed in accordance with the NIH *Guide for the Principles of Animal Care*. The Animal Ethics Committee

at the Institute of Zoology, Chinese Academy of Sciences approved our protocol and study (Permit No: IOZ14001).

**Data accessibility.** Data available from the Dryad Digital Repository: <http://dx.doi.org/10.5061/dryad.hk1vv> [67].

**Authors' contributions.** B.-J.S., R.S., and W.-G.D. designed the study; B.-J.S., T.L., Y.M., and J.-K.M. conducted the experiments. B.-J.S. and W.-G.D. analysed data; B.-J.S., A.G., R.S., and W.-G.D. wrote the manuscript.

**Competing interests.** The authors declare no competing interests.

**Funding.** This work was supported by grants from National Natural Science Fund for Distinguished Young Scholars (31525006), the National Key Research and Development Program of China (2016YFC0503200) and the Australian Research Council.

**Acknowledgements.** We thank T.T.W., X.Z.H., and P.C. for their assistance in the laboratory. We are grateful to two anonymous reviewers for their suggestions.

## References

- Lande R. 1980 Sexual dimorphism, sexual selection, and adaptation in polygenic characters. *Evolution* **34**, 292–305. (doi:10.2307/2407393)
- Shine R. 1989 Ecological causes for the evolution of sexual dimorphism: a review of the evidence. *Quart. Rev. Biol.* **64**, 419–461. (doi:10.1086/416458)
- Breedlove SM. 1992 Sexual dimorphism in the vertebrate nervous-system. *J. Neuro.* **12**, 4133–4142.
- West SA, Reece SE, Sheldon BC. 2002 Sex ratios. *Heredity* **88**, 117–124. (doi:10.1038/sj.hdy.6800018)
- Sarre SD, Georges A, Quinn A. 2004 The ends of a continuum: genetic and temperature-dependent sex determination in reptiles. *BioEssays* **26**, 639–645. (doi:10.1002/bies.20050)
- Kokko H, Jennions MD. 2008 Parental investment, sexual selection and sex ratios. *J. Evol. Biol.* **21**, 919–948. (doi:10.1111/j.1420-9101.2008.01540.x)
- Lang JW, Andrews HV. 1994 Temperature-dependent sex determination in crocodylians. *J. Exp. Zool.* **270**, 28–44. (doi:10.1002/jez.1402700105)
- Ewert MA, Nelson CE. 1991 Sex determination in turtles: diverse patterns and some possible adaptive values. *Copeia* **1991**, 50–69. (doi:10.2307/1446248)
- Harlow P. 2004 Temperature dependent sex determination in lizards. In *Temperature dependent sex determination in vertebrates* (eds N Valenzuela, VA Lance), pp. 11–20. Washington, DC: Smithsonian Institution.
- Cree A, Thompson MB, Daugherty CH. 1995 Tuatara sex determination. *Nature* **375**, 543. (doi:10.1038/375543a0)
- Charnov EL, Bull JJ. 1977 When is sex environmentally determined? *Nature* **266**, 828–830. (doi:10.1038/266828a0)
- Shine R. 1999 Why is sex determined by nest temperature in many reptiles? *Trends Ecol. Evol.* **14**, 186–189. (doi:10.1016/S0169-5347(98)01575-4)
- Warner DA, Shine R. 2008 The adaptive significance of temperature-dependent sex determination in a reptile. *Nature* **451**, 566–568. (doi:10.1038/nature06519)
- Rhen T, Schroeder A. 2010 Molecular mechanisms of sex determination in reptiles. *Sex Dev.* **4**, 16–28. (doi:10.1159/000282495)
- Matsumoto Y, Crews D. 2012 Molecular mechanisms of temperature-dependent sex determination in the context of ecological developmental biology. *Mol. Cell Endocr.* **354**, 103–110. (doi:10.1016/j.mce.2011.10.012)
- Merchant-Larios H, Diaz-Hernandez V. 2013 Environmental sex determination mechanisms in reptiles. *Sex Dev.* **7**, 95–103. (doi:10.1159/000341936)
- Webb GJW, Smith AMA. 1984 Sex ratio and survivorship in the Australian freshwater crocodile *Crocodylus johnstoni*. *Symp. Zool. Soc.* **52**, 319–355.
- Webb GJW, Beal AM, Manolis SC, Demsey KE. 1987 The effects of incubation temperature on sex determination and embryonic development rate in *Crocodylus johnstoni* and *C. porosus*. In *Wildlife management of crocodiles and alligators* (eds JW Webb, SC Manolis, PJ Whitehead), pp. 507–531. Sydney, Australia: Surey Beatty and Sons.
- Packard GC, Packard MJ, Benigan L. 1991 Sexual differentiation, growth, and hatching success by embryonic painted turtles incubated in wet and dry environments at fluctuating temperatures. *Herpetologica* **47**, 125–132.
- Etchberger CR, Phillips JB, Ewert MA, Nelson CE, Prange HD. 1991 Effects of oxygen concentration and clutch on sex determination and physiology in red-eared slider turtles (*Trachemys scripta*). *J. Exp. Zool.* **258**, 394–403. (doi:10.1002/jez.1402580315)
- Lockwood SF, Holland BS, Bickham JW, Hanks BG, Bull JJ. 1991 Intraspecific genome size variation in a turtle (*Trachemys scripta*) exhibiting temperature-dependent sex determination. *Can. J. Zool.* **69**, 2306–2310. (doi:10.1139/z91-324)
- Crews D. 1996 Temperature-dependent sex determination: the interplay of steroid hormones and temperature. *Zool. Sci.* **13**, 1–13. (doi:10.2108/zsj.13.1)
- Crews D, Bergeron JM. 1994 Role of reductase and aromatase in sex determination in the red-eared slider (*Trachemys scripta*), a turtle with temperature-dependent sex determination. *J. Endocr.* **143**, 279–289. (doi:10.1677/joe.0.1430279)
- Matsumoto Y, Yatsu R, Taylor C, Crews D. 2013 Changes in gonadal gene network by exogenous ligands in temperature-dependent sex determination. *J. Mol. Endocr.* **50**, 389–400. (doi:10.1530/jme-12-0260)
- Crews D, Cantu AR, Rhen T, Vohra R. 1996 The relative effectiveness of estrone, estradiol-17 beta, and estriol in sex reversal in the red-eared slider (*Trachemys scripta*), a turtle with temperature-dependent sex determination. *Gen. Comp. Endocr.* **102**, 317–326. (doi:10.1006/gcen.1996.0075)
- Rhen T, Lang JW. 1999 Incubation temperature and sex affect mass and energy reserves of hatchling snapping turtles, *Chelydra serpentina*. *Oikos* **86**, 311–319. (doi:10.2307/3546448)
- Carr JA, Patino R. 2011 The hypothalamus-pituitary-thyroid axis in teleosts and amphibians: endocrine disruption and its consequences to natural populations. *Gen. Comp. Endocr.* **170**, 299–312. (doi:10.1016/j.ygcen.2010.06.001)
- Hulbert AJ. 2000 Thyroid hormones and their effects: a new perspective. *Biol. Rev.* **75**, 519–631. (doi:10.1017/s146479310000556x)
- McNabb FMA. 2006 Avian thyroid development and adaptive plasticity. *Gen. Comp. Endocr.* **147**, 93–101. (doi:10.1016/j.ygcen.2005.12.011)
- Walpita CN, Van der Geyten S, Rurangwa E, Darras VM. 2007 The effect of 3,5,3'-triiodothyronine supplementation on zebrafish (*Danio rerio*) embryonic development and expression of iodothyronine deiodinases and thyroid hormone receptors. *Gen. Comp. Endocr.* **152**, 206–214. (doi:10.1016/j.ygcen.2007.02.020)
- Power DM, Llewellyn L, Faustino M, Nowell MA, Bjornsson BT, Einarsdottir IE, Canario AVM, Sweeney GE. 2001 Thyroid hormones in growth and development of fish. *Comp. Biochem. Physiol. C* **130**, 447–459. (doi:10.1016/s1532-0456(01)00271-x)
- Shepherdley CA, Richardson SJ, Evans BK, Kuhn ER, Darras VM. 2002 Thyroid hormone deiodinases during embryonic development of the saltwater crocodile (*Crocodylus porosus*). *Gen. Comp. Endocr.* **126**, 153–164. (doi:10.1006/gcen.2002.7786)
- De Groef B, Grommen SVH, Darras VM. 2013 Hatching the cleidoic egg: the role of thyroid hormones. *Front. Endocr.* **4**, 63. (doi:10.3389/fendo.2013.00063)
- Hayes TB. 1998 Sex determination and primary sex differentiation in amphibians: genetic and developmental mechanisms. *J. Exp. Zool.* **281**, 373–399. (doi:10.1002/(sici)1097-010x(19980801)281:5<373::aid-jez4>3.3.co;2-t)
- Goleman WL, Carr JA, Anderson TA. 2002 Environmentally relevant concentrations of ammonium perchlorate inhibit thyroid function and alter sex ratios in developing *Xenopus laevis*. *Environ. Toxicol. Chem.* **21**, 590–597. (doi:10.1897/1551-5028(2002)021<0590:ercoap>2.0.co;2)
- Mukhi S, Torres L, Patino R. 2007 Effects of larval-juvenile treatment with perchlorate and

- co-treatment with thyroxine on zebrafish sex ratios. *Gen. Comp. Endocr.* **150**, 486–494. (doi:10.1016/j.ygcen.2006.11.013)
37. Wibbels T, Cowan J, LeBoeuf R. 1998 Temperature-dependent sex determination in the red-eared slider turtle, *Trachemys scripta*. *J. Exp. Zool.* **281**, 409–416. (doi:10.1002/(sici)1097-010x(19980801)281:5<409::aid-jez6>3.0.co;2-s)
38. Ernst CH, Lovich JE. 2009 *Turtles of the United States and Canada*. Baltimore, MA: Johns Hopkins University Press.
39. Godfrey MH, Delmas V, Girondot M. 2003 Assessment of patterns of temperature-dependent sex determination using maximum likelihood model selection. *Ecoscience* **10**, 265–272.
40. Jackson DR. 1994 Overwintering of hatchling turtles in northern Florida. *J. Herpetol.* **28**, 401–402. (doi:10.2307/1564549)
41. Sun BJ, Li T, Gao J, Ma L, Du WG. 2015 High incubation temperatures enhance mitochondrial energy metabolism in reptile embryos. *Sci. Rep.* **5**, 8861. (doi:10.1038/Srep08861)
42. O'Steen S, Janzen FJ. 1999 Embryonic temperature affects metabolic compensation and thyroid hormones in hatchling snapping turtles. *Physiol. Biochem. Zool.* **72**, 520–533. (doi:10.1086/316690)
43. McGlashan JK, Thompson MB, Van Dyke J, Spencer RJ. In press. Thyroid hormones reduce incubation period without developmental and metabolic costs in Murray River short-necked turtles (*Emydura macquarii*). *Physiol. Biochem. Zool.*
44. Yntema CL. 1968 A series of stages in embryonic development of *Chelydra serpentina*. *J. Morphol.* **125**, 219–251. (doi:10.1002/jmor.1051250207)
45. Greenbaum E. 2002 A standardized series of embryonic stages for the emydid turtle *Trachemys scripta*. *Can. J. Zool.* **80**, 1350–1370. (doi:10.1139/z02-111)
46. Rhen T, Fagerlie R, Schroeder A, Crossley DA, Lang JW. 2015 Molecular and morphological differentiation of testes and ovaries in relation to the thermosensitive period of gonad development in the snapping turtle, *Chelydra serpentina*. *Differentiation* **89**, 31–41. (doi:10.1016/j.diff.2014.12.007)
47. Ramsey M, Shoemaker C, Crews D. 2007 Gonadal expression of Sf1 and aromatase during sex determination in the red-eared slider turtle (*Trachemys scripta*), a reptile with temperature-dependent sex determination. *Differentiation* **75**, 978–991. (doi:10.1111/j.1432-0436.2007.00182.x)
48. Du WG, Radder RS, Sun B, Shine R. 2009 Determinants of incubation period: do reptilian embryos hatch after a fixed total number of heart beats? *J. Exp. Biol.* **212**, 1302–1306. (doi:10.1242/jeb.027425)
49. McGlashan JK, Loudon FK, Thompson MB, Spencer R-J. 2015 Hatching behavior of eastern long-necked turtles (*Chelodina longicollis*): the influence of asynchronous environments on embryonic heart rate and phenotype. *Comp. Biochem. Physiol. A* **188**, 58–64. (doi:10.1016/j.cbpa.2015.06.018)
50. Gutzke WHN, Paukstis GL, Packard GC. 1984 Pipping versus hatching as indexes of time of incubation in reptiles. *J. Herpetol.* **18**, 494–496. (doi:10.2307/1564114)
51. Du WG, Hu LJ, Lu JL, Zhu LJ. 2007 Effects of incubation temperature on embryonic development rate, sex ratio and post-hatching growth in the Chinese three-keeled pond turtle, *Chinemys reevesii*. *Aquaculture* **272**, 747–753. (doi:10.1016/j.aquaculture.2007.09.009)
52. Denver RJ, Licht P. 1988 Thyroid status influences *in vitro* thyrotropin and growth-hormone responses to thyrotropin-releasing-hormone by pituitary-glands of hatchling slider turtles (*Pseudemys scripta elegans*). *J. Exp. Zool.* **246**, 293–304. (doi:10.1002/jez.1402460309)
53. Forhead AJ, Fowden AL. 2014 Thyroid hormones in fetal growth and parturition maturation. *J. Endocr.* **221**, R87–R103. (doi:10.1530/joe-14-0025)
54. Yen PM. 2001 Physiological and molecular basis of thyroid hormone action. *Physiol. Rev.* **81**, 1097–1142.
55. Sharma P, Patiño R. 2013 Regulation of gonadal sex ratios and pubertal development by the thyroid endocrine system in zebrafish (*Danio rerio*). *Gen. Comp. Endocr.* **184**, 111–119. (doi:10.1016/j.ygcen.2012.12.018)
56. Georges A, Limpus C, Stoutjesdijk R. 1994 Hatching sex in the marine turtle *Caretta caretta* is determined by proportion of development at a temperature, not daily duration of exposure. *J. Exp. Zool.* **270**, 432–444. (doi:10.1002/jez.1402700504)
57. Du WG, Shen JW, Wang L. 2009 Embryonic development rate and hatchling phenotypes in the Chinese three-keeled pond turtle (*Chinemys reevesii*): the influence of fluctuating temperature versus constant temperature. *J. Therm. Biol.* **34**, 250–255. (doi:10.1016/j.jtherbio.2009.03.002)
58. Neuwald JL, Valenzuela N. 2011 The lesser known challenge of climate change: thermal variance and sex-reversal in vertebrates with temperature-dependent sex determination. *PLoS ONE* **6**, e18117. (doi:10.1371/journal.pone.0018117)
59. Schmidt F, Schnurr S, Wolf R, Braunbeck T. 2012 Effects of the anti-thyroidal compound potassium-perchlorate on the thyroid system of the zebrafish. *Aqua Toxicol.* **109**, 47–58. (doi:10.1016/j.aquatox.2011.11.004)
60. Bernhardt RR, von Hippel FA, Cresko WA. 2006 Perchlorate induces hermaphroditism in threespine sticklebacks. *Environ. Toxicol. Chem.* **25**, 2087–2096. (doi:10.1897/05-454r.1)
61. Vasudevan N, Ogawa S, Pfaff D. 2002 Estrogen and thyroid hormone receptor interactions: physiological flexibility by molecular specificity. *Physiol. Rev.* **82**, 923–944. (doi:10.1152/physrev.00014.2002)
62. Uchida D, Yamashita M, Kitano T, Iguchi T. 2004 An aromatase inhibitor or high water temperature induce oocyte apoptosis and depletion of P450 aromatase activity in the gonads of genetic female zebrafish during sex-reversal. *Comp. Biochem. Physiol. A* **137**, 11–20. (doi:10.1016/s1095-6433(03)00178-8)
63. Fazli N, Hassanabadi A, Mottaghitalab M, Hajati H. 2015 Manipulation of broiler chickens sex differentiation by *in ovo* injection of aromatase inhibitors, and garlic and tomato extracts. *Poult. Sci.* **94**, 2778–2783. (doi:10.3382/ps/pev236)
64. Arambepola NK, Bunick D, Cooke PS. 1998 Thyroid hormone effects on androgen receptor messenger RNA expression in rat Sertoli and peritubular cells. *J. Endocr.* **156**, 43–50. (doi:10.1677/joe.0.1560043)
65. Cardone A, Angelini F, Esposito T, Comitato R, Varriale B. 2000 The expression of androgen receptor messenger RNA is regulated by tri-iodothyronine in lizard testis. *J. Ster. Biochem. Mol. Biol.* **72**, 133–141. (doi:10.1016/s0960-0760(00) 00021-2)
66. Valenzuela N, Lance V. 2004 *Temperature dependent sex determination in vertebrates*. Washington, DC: Smithsonian Books.
67. Sun BJ, Li T, Mu Y, McGlashan J, Georges A, Shine R, Du WG. 2016 Data from: thyroid hormone modulates offspring sex ratio in a turtle with temperature-dependent sex determination. Dryad Digital Repository. (doi:10.5061/dryad.hk1v)